

Chiral medium produced by parallel electric and magnetic fields

Marco Ruggieri^{1,*}, Guang Xiong Peng^{1,2,**}, and Maxim Chernodub^{3,4,***}

¹College of Physics, University of Chinese Academy of Sciences, Yuquanlu 19A, Beijing 100049, China

²Theoretical Physics Center for Science Facilities, Institute of High Energy Physics, Beijing 100049, China

³Laboratoire de Mathématiques et Physique Théorique UMR 7350, Université de Tours, 37200, France

⁴Laboratory of Physics of Living Matter, Far Eastern Federal University, Sukhanova 8, Vladivostok, Russia

Abstract. We compute (pseudo)critical temperature, T_c , of chiral symmetry restoration for quark matter in the background of parallel electric and magnetic fields. This field configuration leads to the production of a chiral medium on a time scale τ , characterized by a nonvanishing value of the chiral density that equilibrates due to microscopic processes in the thermal bath. We estimate the relaxation time τ to be about $\approx 0.1 - 1$ fm/c around the chiral crossover; then we compute the effect of the fields and of the chiral medium on T_c . We find T_c to be lowered by the external fields in the chiral medium.

1 Introduction

In recent years it has been realized that in ultrarelativistic heavy ion collisions very strong electric and magnetic fields can be produced, at least in the very early stages of the collision [1, 2]. In particular the strong magnetic field and the interplay with chiral (ABJ) anomaly in Quantum Chromodynamics (QCD) [3, 4] has suggested the possibility of observation of chiral magnetic effect (CME) in relativistic heavy ion collisions [5, 6]; a recent review and an exhaustive list of references can be found in [7]. Naming $n_{R/L}$ the number densities of right- and left-handed species, and \mathcal{E} , \mathcal{B} a color-electric and color-magnetic field respectively, CME is related to the generation of chiral density, $n_5 = n_R - n_L$ induced by interaction of quarks with gluon fields characterized by $\mathcal{E} \cdot \mathcal{B} \neq 0$ named the QCD sphalerons.

Because of chiral anomaly it is possible to induce a chiral density also by the interaction of quark matter with external electric, E , and magnetic, B , fields with $E \cdot B \neq 0$. The chiral anomaly will then inject n_5 into the system, and this chiral density can equilibrate thanks to microscopic processes that change quark chirality. The purpose of the study presented here is to study this equilibration, which leads to the formation of a medium that we name chiral medium, and to compute its effect on the critical temperature of QCD. The work presented here is mainly based on [8–12].

**Speaker.* e-mail: marco.ruggieri@ucas.ac.cn

**e-mail: gxpeng@ucas.ac.cn

***e-mail: maxim.chernodub@lmpt.univ-tours.fr

2 Chiral medium produced by E and B

We consider quark matter in the background of parallel Abelian (electromagnetic) electric, E , and magnetic, B , fields [8–10] (see also [13–15]). In this case the evolution of n_5 with time is given by

$$\frac{dn_5}{dt} = -n_5\Gamma + N_c \frac{e^2 E \cdot B}{2\pi^2} \sum_f q_f^2 e^{-\frac{\pi M_q^2}{|q_f e E|}}, \quad (1)$$

where the first term on the right hand side describes relaxation of chiral density due to chirality-changing processes in the thermal medium, while the second term comes from the ABJ chiral anomaly [3, 4, 13]. M_q denotes the quark mass and q_f the electric charge of the flavor f in units of the proton charge. The quantity Γ in Eq. (1) corresponds to the rate of the chirality flips while its inverse defines the chiral relaxation time,

$$\tau = 1/\Gamma. \quad (2)$$

The physical interpretation of Eq. (1) is that in the background of $E \parallel B$ fields the ABJ anomaly creates a chiral imbalance n_5 ; the growth of n_5 competes however with chiral relaxation due to microscopic processes that flip the chirality of quarks. For time $t \approx \tau$ the value of chiral density exponentially equilibrates at:

$$n_5^{\text{eq}} = N_c \frac{e^2 E \cdot B}{2\pi^2} \tau \sum_f q_f^2 e^{-\frac{\pi M_q^2}{|q_f e E|}}. \quad (3)$$

The main message encoded in Eq. (3) is that the combined effect of the external fields and of the chirality changing processes in the thermal bath is the formation of chiral medium, characterized by a finite value of chiral density and to which a chiral chemical potential μ_5 can be associated, namely

$$n_5^{\text{eq}} = -\frac{\partial \Omega}{\partial \mu_5}, \quad (4)$$

where Ω corresponds to the thermodynamic potential.

The knowledge of the relaxation time τ is crucial since it allows to determine the equilibrium value of chiral density, n_5^{eq} . In the context of QCD, τ been computed in [9] for temperatures around T_c within an NJL model. The microscopic processes considered in [9] are quark-quark scattering mediated by collective modes with the quantum numbers of pions and σ -mesons, hence we refer to these processes as one-pion or one- σ exchange for simplicity. While referring the interested reader to [9] for all the technical details, we remind here the main quantities that have to be computed to estimate τ . The starting point is the Boltzmann collision integral,

$$\frac{df_R(p)}{dt} = \int d\Pi \frac{(2\pi)^4 \delta^4(p+k-p'-k')}{2E_p} |\mathcal{M}|^2 F, \quad (5)$$

where f_R denotes the distribution function for the right-handed quarks (a similar equation holds for left-handed quarks) and $d\Pi$ corresponds to the momentum space measure,

$$d\Pi = \frac{d^3k}{(2\pi)^3 2E_k} \frac{d^3k'}{(2\pi)^3 2E'_k} \frac{d^3p'}{(2\pi)^3 2E'_p}; \quad (6)$$

the kernel F takes into account the population of the incoming and outgoing particles in the process,

$$F(p, k, p', k') = f_L(p') f_L(k') [1 - f_R(p)] [1 - f_R(k)] - f_R(p) f_R(k) [1 - f_L(p')] [1 - f_L(k')], \quad (7)$$




Figure 1. Left panel: relaxation time versus temperature in the crossover region for π -exchange (squares) and σ -exchange (circles). Right panel: relaxation time for π -exchange for the cases of the pure NJL model (green diamonds) and of the NJL+quasiparticle model (blue triangles). Adapted from [9].

for the scattering of two incoming R quarks giving two outgoing L quarks, with $f_{R/L}(p) = (1 + e^{\beta\omega_{\pm}})^{-1}$ and $\omega_s = \sqrt{(p + s\mu_5)^2 + M_q^2}$ with $s = \pm 1$. In [9] any dependence of τ on E and B has been neglected for simplicity, and an expansion to the lowest nontrivial order in μ_5/T has been used to further simplify the numerical evaluation of the collision integral.

In Eq.(5) the squared transition amplitude $|\mathcal{M}|^2$ can be computed once a microscopic process for the chirality change has been chosen, namely the π -exchange and σ -exchange in [9]. From the definition of n_5 we get the equation governing its time evolution, namely

$$\frac{dn_5}{dt} = N_c N_f \int \frac{d^3p}{(2\pi)^3} \left(\frac{df_R}{dt} - \frac{df_L}{dt} \right). \quad (8)$$

Then from Eq.(1) with $E = B = 0$ we get the collision rate $n_5\Gamma = -dn_5/dt$; the relation between n_5 and μ_5 has been computed in [9] within the NJL model. Once the reaction rate Γ is known the relaxation time can be computed by Eq.(2).

In the left panel of Fig. 1 we plot the relaxation time versus temperature in the chiral crossover region for π -exchange (squares) and σ -exchange (circles). We notice that the σ -exchange is less efficient than π -exchange in changing chirality of quarks in the thermal bath, resulting in a larger relaxation time. This can be understood because of the larger thermal mass of the σ -meson with respect to the pion mass.

An interesting feature of the results shown in Fig. 1 is that the relaxation time decreases with temperature. This feature might sound counterintuitive: naively a chirality change in the system should be related to the quark mass operator, that in the NJL model is related to M_q which becomes smaller with temperature, implying Γ becomes also smaller. However, while this naive argument works well for the transition matrix element, it does not necessarily work for Γ : in fact the latter is determined also by the phase space available for collisions, which instead becomes larger with temperature because the quark mass becomes smaller and the distribution functions broader. As a result, the collision rate increases with temperature and the relaxation time becomes smaller. For a more detailed discussion see [9].

In [9] the NJL model calculation of τ has been compared with a model calculation where the quark mass function at large temperature is taken to be that of a quasiparticle model [16]. The latter is




Figure 2. Left panel: Critical temperature versus μ_5 . Data taken from [11]. Right panel: M_q versus temperature for the case $eE = eB = 8m_\pi^2$ and $n_5 = 0$ (green dots), same values of E and B but with n_5 given by its equilibrium value Eq. (3) (orange dashed line). For comparison we plot by indigo solid line the data corresponding to $E = B = 0$. Adapted from [10].

assumed to take into account many-body effects that are ignored within the NJL model: in particular, the quasiparticle mass is larger than temperature, at least for $T \approx T_c$, while the NJL mass becomes much smaller than the temperature itself because of chiral symmetry restoration. The smooth transition from the NJL mass at low temperature to the quasiparticle mass at large temperature introduced in [9] models a phase space volume reduction at $T \approx T_c$, hence potentially lowering Γ and increasing τ at the crossover. In the right panel of Fig. 1 we plot the relaxation time for π -exchange computed within the NJL model (green diamonds) and the NJL+quasiparticle model (blue triangles). The main difference of the relaxation time computed within the NJL+quasiparticle model in [9], with respect to the one obtained within the pure NJL model, is that in the former case τ stays almost constant at the crossover instead of decreasing as in the pure NJL model calculation; moreover, its order of magnitude is $\tau \approx 1$ fm/c while for the NJL calculation it drops down to $\tau \approx 0.1$ fm/c for $T > T_c$. Clearly the merging of NJL and quasiparticle models of [9] is too naive: nevertheless the results about τ show that the many body interactions, usually neglected in NJL model calculations, can have important effects on the computation of physical quantities around and above T_c , and more serious studies about these effects should be performed.

3 Critical temperature for chiral symmetry restoration

A natural question that arises is: can the formation of the chiral medium lead to any observable effect on equilibrium properties of the bulk? In particular, in our study we are interested to the QCD critical temperature in presence of the electric and magnetic fields, therefore we can ask whether the equilibrated n_5 can affect T_c . The answer is that in principle this is possible. To elaborate more on this point we plot in the left panel of Fig. 2 the critical temperature versus μ_5 computed in [11] within NJL models with several interaction kernels for $E = B = 0$. The results in Fig. 2 show that T_c increases with μ_5 : the chiral chemical potential acts as a catalyzer of chiral symmetry breaking, see also [17–20]. Since in the context of our study the chemical potential is produced by the external fields, the net effect of the field on T_c has to be considered taking into account the equilibrated n_5 .

The effect of μ_5 on T_c is opposite to the one induced by the fields. In fact on the right panel of Fig. 2 we plot the constituent quark mass versus temperature for $eE = eB = 8m_\pi^2$, for the cases in




Figure 3. Critical temperature versus electric field strength for several values of B . Adapted from [10].

which we ignore the chiral density at equilibrium (green squares). We also plot the data corresponding to $E = B = 0$ by the solid indigo line. Data have been obtained from [10]. The results collected in the right panel of Fig. 2 show that the fields lower both M_q and T_c in comparison with the $E = B = 0$ case: the fields thus act as inhibitors of spontaneous chiral symmetry breaking, see also [14].

Given the different response of T_c to the fields on the one hand, and to μ_5 on the other hand, it is important to consider both of the effects to make a firm computation of T_c in presence of $E \parallel B$. This program has been accomplished in [10] where the QCD critical temperature in the background of $E \parallel B$ has been computed taking into account the equilibration of chiral density. In the right panel of Fig. 2 we plot by orange dashed line M_q versus temperature for $eE = eB = 8m_\pi^2$ for the case in which the equilibrated chiral density has been taken into account, by solving simultaneously the gap and the number equations within the NJL model. As expected, the effect of the chiral medium is to increase the value of M_q with respect to the $n_5 = 0$ case, as shown by the comparison of the orange and the green lines. On the other hand, the net effect of the chiral medium is not so strong, since the increase of M_q due to $n_5 \neq 0$ is always within the 10% of the $n_5 = 0$ result; as a consequence we conclude that the effect of $E \parallel B$ is much stronger than the one induced by n_5 and the net effect of the fields will be a lowering of T_c . Similar results have been recently obtained in [8].

In [10] the picture drawn here has been completed by the computation of the bulk structure of the chiral medium at equilibrium: in particular, it has been found that the chiral chemical potentials for u and d quarks are different, a result that can be easily understood because n_5 is produced by the electromagnetic coupling of quarks to E and B so a dependence on the quark electric charge is expected. Hence the external fields induce an imbalanced population of u and d quarks, which might have some effect on the charged pion condensation.

In Fig. 3, adapted from [10], we plot the critical temperature as a function of E for several values of B . The general trend of the results shown in the figure is that the electric field lowers T_c . An interesting feature of data is that even in the case of the largest magnetic field considered in the figure, which increases T_c with respect to the case $B = 0$, a modest electric field $eE \approx 4m_\pi^2$ is enough to bring T_c down to the zero field case. Hence the conclusion is that $E \parallel B$ acts as an inhibitor of chiral symmetry breaking, as long as E is not much smaller than B .

4 Conclusions

In this talk we have discussed our results on chiral symmetry restoration at finite temperature, for quark matter in the presence of parallel electric, E , and magnetic, B , fields. This background leads to the formation of a chiral medium, namely a medium with a finite chiral density at equilibrium, $n_5 = n_R - n_L$, because of the combined effect of the chiral anomaly which pumps n_5 into the system, and chirality changing microscopic processes in the thermal bath that instead suppress the formation of n_5 . The characteristic time scale necessary for the formation of this medium corresponds approximately to the relaxation time of chiral density, τ . We have estimated τ around the chiral crossover of QCD, finding $\tau \approx 0.1 - 1$ fm/c.

Then we have used this value of τ to compute n_5 at equilibrium simultaneously with the in-medium chiral condensate at finite temperature, eventually computing the critical temperature, T_c , as a function of the external field magnitude. Our result is summarized in Fig. 3 where we plot T_c versus the electric field strength for several values of the magnetic field; the critical lines have been computed by taking into account the value of n_5 at equilibrium computed self-consistently by virtue of Eq. (3). We find that the $E \parallel B$ background acts as an inhibitor of chiral symmetry breaking, at least for values of E not too smaller than B .

An interesting follow up of the work presented here might be the inclusion of neutral pion condensation which is known to be induced by the chiral anomaly [15]; also it might be worth studying the charged pion condensation induced by the mismatch of u and d chemical potentials in the chiral medium. We plan to report on these subjects in the near future.

Acknowledgements. M. R and G. X. P. would like to thank the CAS President's International Fellowship Initiative (Grant No. 2015PM008), and the NSFC projects (11135011 and 11575190). M. R. acknowledges discussions with V. Greco, Z. Y. Lu and F. Scardina.

References

- [1] V. Skokov, A. Y. Illarionov and V. Toneev, *Int. J. Mod. Phys. A* **24**, 5925 (2009).
- [2] A. Bzdak and V. Skokov, *Phys. Lett. B* **710**, 171 (2012).
- [3] S. L. Adler, *Phys. Rev.* **177**, 2426 (1969).
- [4] J. S. Bell and R. Jackiw, *Nuovo Cim. A* **60**, 47 (1969).
- [5] D. E. Kharzeev, L. D. McLerran and H. J. Warringa, *Nucl. Phys. A* **803**, 227 (2008).
- [6] K. Fukushima, D. E. Kharzeev and H. J. Warringa, *Phys. Rev. D* **78**, 074033 (2008).
- [7] K. Hattori and X. G. Huang, arXiv:1609.00747 [nucl-th].
- [8] M. Ruggieri, Z. Y. Lu and G. X. Peng, arXiv:1608.08310 [hep-ph].
- [9] M. Ruggieri, G. X. Peng and M. Chernodub, *Phys. Rev. D* **94**, no. 5, 054011 (2016).
- [10] M. Ruggieri and G. X. Peng, *Phys. Rev. D* **93**, no. 9, 094021 (2016).
- [11] M. Ruggieri and G. X. Peng, arXiv:1602.05250 [hep-ph].
- [12] M. Ruggieri and G. X. Peng, arXiv:1602.03651 [hep-ph].
- [13] H. J. Warringa, *Phys. Rev. D* **86**, 085029 (2012).
- [14] A. Y. Babansky, E. V. Gorbar and G. V. Shchepanyuk, *Phys. Lett. B* **419**, 272 (1998).
- [15] G. Cao and X. G. Huang, *Phys. Lett. B* **757**, 1 (2016).
- [16] S. Plumari, W. M. Alberico, V. Greco and C. Ratti, *Phys. Rev. D* **84**, 094004 (2011).
- [17] L. Yu, H. Liu and M. Huang, *Phys. Rev. D* **94**, no. 1, 014026 (2016).
- [18] V. V. Braguta *et al.*, *Phys. Rev. D* **93**, no. 3, 034509 (2016).
- [19] V. V. Braguta *et al.*, *JHEP* **1506**, 094 (2015).
- [20] V. V. Braguta and A. Y. Kotov, *Phys. Rev. D* **93**, no. 10, 105025 (2016).