“ Fundamentals of Thermodynamic Modelling of Materials ”

November 15-19, 2010
INSTN – CEA Saclay, France

PROFESSOR & TOPIC

Suzanna FRIES

ICAMS - Ruhr University, Bochum, Germany

*Extrapolations from
thermodynamic
models*

[03004]

Organized by

Bo SUNDMAN bo.sundman@cea.fr
Constantin MEIS constantin.meis@cea.fr

Extrapolations from Thermodynamic Models

Suzana G. Fries

Interdisciplinary Centre for Advanced Materials Simulations
Ruhr University Bochum, Bochum, Germany

Extrapolations from Thermodynamic Models

Lattice stabilities

Metastable regions

Use in other applications

Extrapolations from Thermodynamic Models

Stefano Baroni

why ab initio

scientific insight stems from the ability to extrapolate our understanding from simple models to a very complex physical world

no extrapolation is possible if our knowledge of the models is

- unfeasible
- irreproducible
- irrelevant
- inaccurate
- biased

15 - 19 novembre • INSTN, Saclay, France

« Fundamentals of Thermodynamic Modelling of Materials »

ICAMS

RUB

Extrapolations from Thermodynamic Models

Liquidus extrapolations for Cr

Bo Sundman

15 - 19 novembre • INSTN, Saclay, France

« Fundamentals of Thermodynamic Modelling of Materials »

ICAMS

RUB

Jean-Marc Joubert

Extrapolations from Thermodynamic Models – Models to train...

Gibbs energy	G	$=$	$G(T, p, N_i)$
entropy	S	$=$	$-\left(\frac{\partial G_m}{\partial T}\right)_{p, N_i}$
enthalpy	H	$=$	$G + TS = G - T\left(\frac{\partial G}{\partial T}\right)_{p, N_i}$
volume	V	$=$	$\left(\frac{\partial G}{\partial p}\right)_{T, N_i}$
Chemical potential of component i	μ_i	$=$	$\left(\frac{\partial G}{\partial N_i}\right)_{T, N_{j \neq i}}$
heat capacity	C_p	$=$	$-T\left(\frac{\partial^2 G}{\partial T^2}\right)_{p, N_i}$
thermal expansion	α	$=$	$\frac{1}{V}\left(\frac{\partial^2 G}{\partial p \partial T}\right)_{N_i}$
isothermal compressibility	κ	$=$	$-\frac{1}{V}\left(\frac{\partial^2 G}{\partial p^2}\right)_{T, N_i}$
bulk modulus	B	$=$	$\frac{1}{\kappa}$

Elastic constants

15 - 19 novembre • INSTN, Saclay, France

« Fundamentals of Thermodynamic Modelling of Materials »

ICAMS

RUB

Extrapolations from Thermodynamic Models

Open field : Models can be improved.

More quantities are modelled, higher the consistency for applications

The general Gibbs energy expression is

$$G_m = {}^{\text{srf}}G_m - T {}^{\text{cfg}}S_m + {}^{\text{E}}G_m + {}^{\text{phys}}G_m$$

- ${}^{\text{srf}}G_m$ is the surface of reference for Gibbs energy
- ${}^{\text{cfg}}S_m$ is the configurational entropy
- ${}^{\text{E}}G_m$ is the excess Gibbs energy
- ${}^{\text{phys}}G_m$ is a physical contribution (magnetic)

$$G_m = G_m^{\text{srf}} - TS_m^{\text{conf}} + G_m^{\text{xs}}$$

The surface of reference for the model is the sum of Gibbs energies of all configurations with atom, i , on each sublattice, s , weighted by the site fractions $y_i^{(s)}$. The σ phase has five sublattices with different numbers of sites and coordinations and can be described as

(Re,W)₄(Re,W)₂(Re,W)₈(Re,W)₈(Re,W)₈, sites 1-5, respectively. This gives $2^5=32$ ordered configurations. Thus, the surface of reference becomes

$$G^{\text{srf}} = p_{ijklm} {}^oG_{ijklm}$$

$$p_{ijklm} = y_i^{(1)} y_j^{(2)} y_k^{(3)} y_l^{(4)} y_m^{(5)}$$

15 - 19 novembre • INSTN, Saclay, France

« Fundamentals of Thermodynamic Modelling of Materials »

ICAMS

RUB

These compound energies, ${}^oG_{ijklm}$, are exactly the same FP energies calculated in the paper by Berne et al.

The only additional term necessary to calculate the Gibbs energy at any temperature is the entropy of mixing which is assumed to be ideal in CEF:

$$S^{conf} = -R \sum_s a^{(s)} \sum_{i=Re,W} y_i^{(s)} \ln(y_i^{(s)})$$

$$\left(\frac{\partial G_m}{\partial T} \right)_{p,x_i} = \left(\frac{\partial G_m}{\partial T} \right)_{p,y_j} + \sum_j \left(\frac{\partial G_m}{\partial y_j} \right)_{T,p,y_k} \cdot \left(\frac{dy_j}{dT} \right)_{p,y_k} \quad (k \neq j) \quad (5)$$

The second derivative of the Gibbs energy with respect to mole fractions is

$$\Omega_{ij} = \left(\frac{\partial^2 G}{\partial x_i \partial x_j} \right)_{T,p,x_k} \quad (k \neq i, k \neq j) \quad (6)$$

Extrapolations from Thermodynamic Models

Extrapolations from Thermodynamic Models

Ag-Sn-Cu

15 - 19 novembre • INSTN, Saclay, France

« Fundamentals of Thermodynamic Modelling of Materials »

ICAMS

RUB

Extrapolations from Thermodynamic Models

:5, N=1, T=673;

THERMO-CALC (2009.12.14:09.12) :CU SN
DATABASE TCBN
P=1E5, N=1, T=673;

15 - 19 novembre • INSTN, Saclay, France

« Fundamentals of Thermodynamic Modelling of Materials »

ICAMS

RUB

Extrapolations from Thermodynamic Models

Ag-Sn-Cu

15 - 19 novembre • INSTN, Saclay, France

« Fundamentals of Thermodynamic Modelling of Materials »

ICAMS

RUB

Extrapolations from Thermodynamic Models

Ag-Sn-Cu

15 - 19 novembre • INSTN, Saclay, France

« Fundamentals of Thermodynamic Modelling of Materials »

ICAMS

RUB

Assessments evolution

Calphad modern Calphad ++	Massalsky	Experimental Phase diagram	Zero th Critical evaluation	Drawn phase diagrams	

Al-Ti

Sundman et al.

Mark Asta, CVM

2007, Sundman et al

Re-W: only VASP total energies and a Bragg-Williams ideal entropy

Calculated diagram using CEF

PHASE CHI % 41 4 12 12!
 CONSTITUENT CHI :RE,W : RE,W : RE,W : RE,W : !
 PARAMETER G(CHI,RE:RE:RE:RE:0) 2.98150E+02 +.079453*SF29#;
 6.00000E+03 N REF0!
 PARAMETER G(CHI,W:RE:RE:RE:0) 2.98150E+02 +.065794*SF29#; 6.00000E+03
 N REF0!
 PARAMETER G(CHI,W:W:RE:RE:0) 2.98150E+02 +.029908*SF29#; 6.00000E+03
 N REF0!
 PARAMETER G(CHI,RE:W:RE:RE:0) 2.98150E+02 +.131924*SF29#; 6.00000E+03
 N REF0!
 PARAMETER G(CHI,W:RE:W:RE:0) 2.98150E+02 +.117119*SF29#; 6.00000E+03
 N REF0!
 PARAMETER G(CHI,RE:W:W:RE:0) 2.98150E+02 +.085917*SF29#; 6.00000E+03
 N REF0!
 PARAMETER G(CHI,W:W:W:RE:0) 2.98150E+02 +.079211*SF29#; 6.00000E+03
 N REF0!
 PARAMETER G(CHI,RE:RE:RE:W:0) 2.98150E+02 +.246515*SF29#; 6.00000E+03
 N REF0!
 PARAMETER G(CHI,W:RE:RE:W:0) 2.98150E+02 +.227301*SF29#; 6.00000E+03
 N REF0!
 PARAMETER G(CHI,RE:W:RE:W:0) 2.98150E+02 +.175929*SF29#; 6.00000E+03
 N REF0!
 PARAMETER G(CHI,W:W:RE:W:0) 2.98150E+02 +.160246*SF29#; 6.00000E+03
 N REF0!
 PARAMETER G(CHI,RE:RE:W:W:0) 2.98150E+02 +.35128*SF29#; 6.00000E+03
 N REF0!
 PARAMETER G(CHI,W:RE:W:W:0) 2.98150E+02 +.335662*SF29#; 6.00000E+03
 N REF0!
 PARAMETER G(CHI,RE:W:W:W:0) 2.98150E+02 +.315773*SF29#; 6.00000E+03
 N REF0!
 PARAMETER G(CHI,W:W:W:W:0) 2.98150E+02 +.304247*SF29#; 6.00000E+03
 N REF0!

Dft energies from
 T. Hammerschmidt et al,

CALPHAD assessment,

On going work by Arthur Bialon

$$G^\sigma = H_{ref}^\sigma - TS_{conf}^\sigma$$

$$H_{ref}^\sigma = \sum_{ijklm} y_i^{(1)} y_j^{(2)} y_l^{(3)} y_m^{(4)} y_n^{(5)} H_{ijklm}$$

$$G^\sigma = \sum_i y_i G_i^\sigma + \sum_{i,j} y_i y_j G_{ij}^\sigma + \sum_{i,j,k} y_i y_j y_k G_{ijk}^\sigma + \dots$$

Computational Thermodynamics

What is it CALPHAD?

It is **not** a software!

It is a method!

Gibbs energies for all the stable phases of a system are described by parametric models. These parameters are obtained in such a way that the Gibbs energy and its derivatives reproduce the experimental corresponding value at its best.

When this is done, by Gibbs energy minimization the phase diagram can be calculated!
CALculated PHase Diagrams

Big advantages: interpolations and extrapolations can be done. Less experiments are needed. Even some anticipation of higher order systems can be done!

Interpolations are so good as the description...Extrapolations too and they are not truly predictable.

If there is a ternary compound that is not in the description it can not be guessed by the extrapolation from the binaries.

Computational Thermodynamics

An overview

A puzzle to be solved till the end of the week:
to which lecture is each one of the images related?

15 - 19 novembre • INSTN, Saclay, France

Winter School : « Fundamentals of Thermodynamic Modelling of Materials »