Experiment and Modelling in Structural NMR

November 28th – December 2nd 2011

INSTN – CEA Saclay, France

Gaël de Paëpe
CEA INAC, France

Hyperpolarization
in magnetic resonance
Special focus on
High Field MAS DNP

[05001]

Organized by

Thibault Charpentier
Patrick Berthault
Constantin Meis

thibault.chapentier@cea.fr
patrick.berthault@cea.fr
constantin.meis@cea.fr

Hyperpolarization in Magnetic Resonance

Special focus on
High Field MAS Dynamic Nuclear Polarization

Winter School, Saclay Nov. 28th – Dec. 2nd

Gaël De Paëpe

Direction des Sciences de la Matière

GRENOBLE
UNIVERSITÉ

Disclaimer:

This is not intended to be an exhaustive review on hyper-polarization techniques in magnetic resonance.

The topic is **too large** to be covered here!

This presentation is directly reflecting the author's own interest and is in no way a qualitative/quantitative comparisons between techniques.

GRENOBLE
UNIVERSITÉ

Nuclear Magnetic Resonance

NMR can be used:

→ as a spectroscopy technique:

Structural and dynamical information at the atomic scale!

→ as a microscopy/imaging technique:

Non-invasive technique, convenient for medical diagnostic!

Fields of application: physics, material sciences, biology and medicine

- Main limitations of the technique: Sensitivity, sensitivity and sensitivity!*

*Griffin's lecture

Direction des Sciences de la Matière

Boltzmann Polarization in NMR

NMR: Low energy transitions

Boltzmann distribution

$$P = \frac{n^+ - n^-}{n^+ + n^-} = \tanh\left(\frac{\gamma\hbar B_0}{2kT}\right) \xrightarrow{\text{High Temperature Approximation}} P \sim \frac{\overbrace{\gamma B_0 \hbar}^{\omega_0/2\pi}}{2kT}$$

- Nuclear polarization is typically of the order of tens of ppm!

Direction des Sciences de la Matière

Signal to noise (S/N) in NMR

Mainly inductive detection... so far

$$S \sim \frac{\gamma_e \gamma_d^2 B_0^{\frac{3}{2}}}{T_s} N_s$$

Broadly accepted expression...

Mainly thermal (Johnson) noise

$$N \sim \sqrt{N_s} \sqrt{T_c}$$

$$S/N \sim \frac{\gamma_e \gamma_d^2 B_0^{\frac{3}{2}}}{\sqrt{T_c} T_s} \sqrt{N_s}$$

B_0 : magnetic field; γ_e/γ_d gyromagnetic ratio of the excited/detected spin
 T_s : Sample temperature; T_c : Circuit temperature

- Goal: enhancement of the signal and reduction of the noise!
- FT NMR, High Field magnet, Cryo-probe, INEPT/CP, etc.

Direction des Sciences de la Matière

Current strategy: going to higher magnetic fields

Polarization as a function of B_0 and sample temperature:

$$P \sim \frac{\gamma B_0 \hbar}{2kT}$$

0.007 % ^1H polarization
 @ 1 GHz / 300 K

CRMN, Lyon

- ~ x 10 polarization gain over the last 2 decades thanks to the development of high field magnets

Direction des Sciences de la Matière

Is this enough? – No!

***We are still off by 4 to 5 orders of magnitude on the Polarization!
(compared to a fully polarized system)***

Solution-state NMR

- 3D structure determination of biomolecules
- Small molecules studies:
Metabolomics, molecular dynamics, etc.

Magnetic Resonance Imaging

- anatomical imaging
- functional imaging
- molecular imaging, etc.

Solid-state NMR

- 3D structure determination of insoluble/non crystalline biomolecules
- Materials studies (low gamma nuclei, etc), etc.

Further increase in sensibility is still high desirable...

Direction des Sciences de la Matière

GRENOBLE
UNIVERSITÉ

What can we do?

Hyperpolarization techniques!

- Usually involve transfer from more ordered system (electron, pH_2 , photon)
- In the following we will concentrate on DNP experiments that are compatible with high field NMR detection!

Direction des Sciences de la Matière

GRENOBLE
UNIVERSITÉ

Hyperpolarization techniques

Since the beginning of NMR, hyperpolarization techniques have always been around...

96,99

- Microwave Induced Dynamic Nuclear Polarization – (mw)-DNP

53,58

87

- Laser-polarized Xenon

75

- Para-hydrogen induced polarization – PHIP

73

- Optically Pumped Dynamic Nuclear Polarization – OPDNP, MI-ODNP

- Photo Chemically Induced Dynamic Nuclear Polarization – Photo-CIDNP

- Haupt effect

- etc.

Direction des Sciences de la Matière

Parahydrogen induced polarization

Direction des Sciences de la Matière

PHIP principles

- Hydrogenation reaction induces an overpopulation of states $\alpha\beta$ and $\beta\alpha$
- Note: the para to ortho conversion is a very slow process even at room temperature!

C. R. Bowers and D. P. Weitekamp, JACS 109 (18), 5541 (1987).

Direction des Sciences de la Matière

PHIP sensitivity gain

AB system – J-coupled

- Up to **five orders** of magnitude of sensitivity enhancement ... in **very short time** ! (seconds to minutes) Moreover the mechanism is **independent of the B_0** .
- **Drawbacks:** the enhanced signal disappears if AB spins are not J-coupled
the signal needs to be detected in high-resolution magnet!
it is necessary to use unsaturated precursor of the target molecule

C. R. Bowers and D. P. Weitekamp, JACS 109 (18), 5541 (1987).

Direction des Sciences de la Matière

Current challenges for PHIP

- PHIP is particularly interesting for **low field NMR** (but not only!)
- PHIP was used to study **hydrogenation reactions**
- **Application to MRI** was demonstrated in 2001 – Recent PHIP experiments involve the use of ^{13}C labeled (metabolically relevant) unsaturated precursors
- **Transfer to other nuclei** is difficult but possible! – this implies the design of hydrogenated molecules with long relaxation time to allow polarization transfer to occur.
- **Long time storage of para-hydrogen** (liquid or gas phase) requires very low paramagnetic impurities.
- **Design adequate pulse sequences** to improve the anti-phase detection (caused by the initial longitudinal order)

K. Golman et al., *Magn. Reson. Med.* 46, 1-5 (2001)

C. R. Bowers and D. P. Weitekamp, *JACS* 109 (18), 5541 (1987).

• etc.

Key authors: Bowers, Weitekamp, Golman, Canet, and others

Direction des Sciences de la Matière

Laser-polarized Xenon

Direction des Sciences de la Matière

Laser-polarized Xenon

Direction des Sciences de la Matière

INAC
INSTITUT NANOSCIENCES
ET CRYOGÉNIE

cea

GRENOBLE
UNIVERSITÉ

cnrs

Laser-polarized Xenon

- The metal ion is **polarized by OP**. This circularly polarized laser beam allows the selective depopulation of one of the two magnetic ground states of Rubidium atoms.
- The induced electronic polarization is transferred by **cross-relaxation to xenon** nuclear spins (or other noble gas).

Key authors: Pines et al., Desvaux, Berthault, etc.

G. Navon, A. Pines and co-workers (Science, 271, 1996, 1848)

Direction des Sciences de la Matière

INAC
INSTITUT NANOSCIENCES
ET CRYOGÉNIE

cea

GRENOBLE
UNIVERSITÉ

cnrs

Preparation of laser-polarized Xenon

Experimental apparatus @ CEA-IRAMIS

Left: the lasers and the optics.

Center: the pumping cell and the coils for creating the magnetic field

Right: the electronics and the pumping group

Typical sensitivity gain @ 10 T, $\epsilon \sim 45\,000$!

- Typical production: 1ml in 5 min at 1 atm

Direction des Sciences de la Matière

Hyperpolarized Xenon – main applications

- **He/Xe MRI** → enhanced anatomical imaging (*gas phase*)
- **Xenon NMR** → bio-sensing using laser polarized xenon
- **Polarization transfer from Xenon to other nuclei: SPINOE/CP experiment**
 - Enhancement of ^1H NMR for the cage molecule (*Saclay, 1999*)
 - Surface-enhanced NMR (*Xe adsorption and NOE transfer*)
 - Exploration of protein hydrophobic cavities, **etc.**

Key authors: Pines et al., Desvaux, Berthault, etc.

Direction des Sciences de la Matière

Hyperpolarized Xenon – challenges & perspectives

For Solution NMR:

- Xenon can easily be dissolved in solution but polarization transfer from the highly polarized Xenon to surrounding nuclei is still an unsolved problem: very weak interaction between xenon and other molecules!

For Magnetic Resonance Imaging:

- towards molecular imaging!

i.e. combination of hyperpolarized Xe MRI and biosensing approach*

Frequency-selective laser-polarized ¹²⁹Xe MRI images: proof of principles

* More detail: see NMR group @ IRAMIS

Key authors: Pines et al., Desvaux, Berthault et al., etc.

Direction des Sciences de la Matière

GRENOBLE
UNIVERSITÉ

Microwave induced DNP

Direction des Sciences de la Matière

GRENOBLE
UNIVERSITÉ

Microwave induced Dynamic Nuclear Polarization

DNP mechanisms:

Overhauser effect – *Metal and liquids*

Solid Effect – *solids*

Cross Effect / Thermal Mixing – *solids*

Two types of interactions:

Fermi contact (isotropic)

Dipolar interaction (anisotropic)

Key authors: Overhauser, Carver, Slichter, Abragam, Goldman, Jeffries, Hill, etc.

Direction des Sciences de la Matière

Dynamic Nuclear Polarization: History

Direction des Sciences de la Matière

Overhauser effect in Metals

Polarization of Nuclear Spins in Metals*

T. R. CARVER† AND C. P. SLICHTER
Department of Physics, University of Illinois, Urbana, Illinois
 (Received August 17, 1953)

- ^7Li NMR @ $\omega_0/2\pi = 50 \text{ kHz}$
 (30.3 Gauss)
- EPR @ $\omega_0/2\pi = 84 \text{ MHz}$

A. Overhauser, *Phys. Rev.* (1953)
 Carver and Slichter, *Phys. Rev.* 92 (1953)
Phys. Rev. 102 (1956)

$\epsilon \sim 100$

FIG. 1. Oscilloscope pictures of 50-ke/sec nuclear resonance absorption vs static magnetic field. Field excursion 0.2 gauss. Top line: Li^7 resonance (lost in noise). Middle line: Li^7 resonance enhanced by electron saturation. Bottom line: Proton resonance in glycerin sample.

- Initial prediction of the effect in 53' by Overhauser
- First demonstration of the Overhauser effect in metals in 53'
- Extension of the effect for liquids!

Direction des Sciences de la Matière

Effect of saturating the EPR transitions in a liquid

Let us consider a two spin system composed of one nuclei and one electron spin

- What have we missed?

Hyperfine coupling between n (I spin) and e (S spin):

$$\left\langle H(t) = \frac{\gamma_e \gamma_n \hbar^2}{r^3} (A + B + C + D + E) \right\rangle = 0$$

$$E = \frac{-3}{4} S^+ I^+ \sin^2(\theta) \exp(-2i\phi) \quad F = \frac{-3}{4} S^- I^- \sin^2(\theta) \exp(2i\phi)$$

$$A = S_z I_z (1 - 3 \cos^2(\theta))$$

$$B = \frac{-1}{4} (S^+ I^- + S^- I^+) (1 - 3 \cos^2(\theta))$$

$$C = \frac{-3}{4} (S^+ I_z + S_z I^+) \sin(2\theta) e^{-i\phi}$$

$$D = \frac{-3}{4} (S^- I_z + S_z I^-) \sin(2\theta) e^{i\phi}$$

Direction des Sciences de la Matière

Solution-state DNP: Overhauser effect

$$W_0 = \frac{1}{20} \frac{\gamma_e^2 \gamma_n^2}{r^6} J(\omega_{0e}, \tau_c)$$

$$W_1 = \frac{6}{20} \frac{\gamma_e^2 \gamma_n^2}{r^6} J(\omega_{0e}, \tau_c)$$

$$W_2 = \frac{3}{20} \frac{\gamma_e^2 \gamma_n^2}{r^6} J(\omega_{0e}, \tau_c)$$

J power density function of the autocorrelation function of H_{dip}

- Motional averaging can induce additional (efficient) relaxation pathways: W_0 , W_1 , W_2
- By saturating the EPR lines, we induce a competition between two processes: saturation (W) and dipolar relaxation towards thermal equilibrium (W_0, W_1, W_2)

Direction des Sciences de la Matière

Effect of saturating the EPR transitions

- Competition between relaxation towards Boltzmann equilibrium and saturation of the EPR lines!

- Complete derivation is possible!

Scalar relaxation: $e \sim -\gamma_e/\gamma_n$

Dipolar relaxation: $e \sim \frac{1}{2} \gamma_e/\gamma_n$

Direction des Sciences de la Matière

High magnetic field regime

Complete derivation is possible...

Coupling parameter

$$\rho(\omega_e, \tau_c) = \frac{W_2 - W_0}{W_0 + 2W_1 + W_2}$$

$$\varepsilon = 1 - \rho f s \frac{\gamma_s}{\gamma_I}$$

Leakage factor $f = \frac{W_0 + 2W_1 + W_2}{W_0 + 2W_1 + W_2 + W_n}$

Saturation Factor $s = \frac{\langle S_0 \rangle - \langle S_z \rangle}{\langle S_0 \rangle} = \frac{W}{W + W_e}$

- High Field Overhauser should not be very efficient!

Loening et al.

Direction des Sciences de la Matière

Solid-state DNP: the Solid Effect

Thermal equilibrium, if B neglected

Hyperfine coupling between e (S spin) and n (I spin):

$$H = \frac{\gamma_e \gamma_n \hbar^2}{r^3} (A + B + C + D + E)$$

$$A = S_z I_z (1 - 3 \cos^2(\theta)) \quad B = \frac{-1}{4} (S^+ I^- + S^- I^+) (1 - 3 \cos^2(\theta))$$

$$C = \frac{-3}{4} (S^+ I_z + S_z I^+) \sin(2\theta) e^{-i\phi} \quad D = \frac{-3}{4} (S^- I_z + S_z I^-) \sin(2\theta) e^{i\phi}$$

$$E = \frac{-3}{4} S^+ I^+ \sin^2(\theta) \exp(-2i\phi) \quad F = \frac{-3}{4} S^- I^- \sin^2(\theta) \exp(2i\phi)$$

Key authors: Abragam

Direction des Sciences de la Matière

Solid-state DNP: the Solid Effect

Thermal equilibrium, if B neglected

Thermal equilibrium, if B not neglected

Key authors: Abragam

Secular approximation for electron Zeeman interaction:

$$H = \omega_{0e} S_z + \omega_{0n} I_z + A S_z I_z + B S_z I_x$$

- No efficient relaxation exist for the zero- or double-quantum transitions, saturation of the EPR transitions is probably not a good idea!

- What happen if we perform an EPR irradiation along the so called "forbidden transitions" ω_0 or ω_2 ? – nothing if B is neglected!

- If B is large enough, it can mix state $\alpha\beta$ and $\beta\beta$ as well as $\alpha\alpha$ and $\beta\alpha$. Then the transitions are not so forbidden anymore...

Direction des Sciences de la Matière

The Solid Effect: irradiating the forbidden transitions

$$\varepsilon \sim \left(\frac{\gamma_e}{\gamma_n} \right) \left(\frac{B_{1mw}}{B_0} \right)^2$$

SE DNP enhancement

- Note: $1/B_0^2$ dependency!!

Direction des Sciences de la Matière

The well-resolved solid Effect

$\omega_{n0} > \text{EPR linewidth}$

$\omega_{n0} \sim \text{EPR linewidth}$

- SE efficiency is inversely proportional to the B_0^2
- SE is applicable when the EPR FWHH is smaller than the ω_{n0}
- But this is not generally the case at high magnetic field...

Direction des Sciences de la Matière

INAC
INSTITUT NANOSCIENCES
ET CRYOGÉNIE

cea

GRENOBLE
UNIVERSITÉ

cnrs

Three spin mechanism DNP: two electrons and one nuclei

Fortunately other mechanisms are possible...which involve more than one electron spin...

Key authors: Kessenikh, Manenkov, Hill, Goldman, Abragam, etc

Direction des Sciences de la Matière

INAC
INSTITUT NANOSCIENCES
ET CRYOGÉNIE

cea

GRENOBLE
UNIVERSITÉ

cnrs

Cross Effect and Thermal Mixing DNP

- Three and more spin process have proven very useful, especially at high B_0
- Detailed mechanisms are complicated!

- EPR inhomogeneous breadth: Δ – g anisotropy
- EPR homogeneous linewidth: δ – ee dipolar couplings
- NMR Larmor Frequency: ω_{0n}

Crude (but simple) classification of the various DNP mechanism...

- if $\omega_{0n} \gg \Delta$: Two spin process (Solid Effect)
 $\epsilon \sim 1/B_0^2$
- if $\delta < \omega_{0n} < \Delta$: Three spin process (Cross Effect)
Very efficient when $\omega_{e1} - \omega_{e2} \sim \omega_{0n}$
 $\epsilon \sim 1/B_0$
- if $\omega_{0n} < \delta$: Multiple electron spin process
Thermal Mixing

Hu et al.

Direction des Sciences de la Matière

Challenges of High Field DNP for solution NMR

- All DNP mechanisms presented here appear less favorable at high magnetic fields!
- Combining DNP with high field NMR also imposes several additional constraints:
 - MW heating is a concern (especially for liquids)
 - Difficult to design resonant structure for the microwave irradiation that are compatible with MAS spinning, etc.

Current on-going research areas:

High Field MAS solid-state DNP (*in situ*, LT, 5-15 T, high power source)

Dissolution DNP (*ex situ* DNP at 1K and 3 Tesla, dissolution, NMR/MRI)

Shuttling DNP for liquids (Low Field Overhauser – High Field NMR)

High Field Overhauser for liquids

Direction des Sciences de la Matière

High Field Dynamic Nuclear Polarization

Direction des Sciences de la Matière

High Field MAS solid-state DNP

Direction des Sciences de la Matière

High Field Solid-state MAS DNP – the Griffin approach

High Field Dynamic Nuclear Polarization

Direction des Sciences de la Matière

High Field Solid-state MAS DNP – the Griffin approach

High Field Dynamic Nuclear Polarization

- DNP results from transferring spin polarizations to surrounding nuclei

Direction des Sciences de la Matière

High Field Solid-state MAS DNP – the Griffin approach

High Field Dynamic Nuclear Polarization

- DNP is produced by applying at the sample an irradiation with a frequency close to the electron spin transition

Direction des Sciences de la Matière

High Field Solid-state MAS DNP – the Griffin approach

High Field Dynamic Nuclear Polarization

$$\epsilon_{\text{max}} = \frac{P_{\text{blue}}}{P_{\text{red}}} = 2000 \rightarrow \text{But recycling is possible (5-10 s) which could translate in } 4 \cdot 10^6 \text{ time savings!!}$$

Direction des Sciences de la Matière

DNP with gyrotrons

VOLUME 71, NUMBER 21

PHYSICAL REVIEW LETTERS

22 NOVEMBER 1993

Dynamic Nuclear Polarization with a Cyclotron Resonance Maser at 5 T

Lino R. Becerra,¹ Gary J. Gerfen,¹ Richard J. Temkin,² David J. Singel,³ and Robert G. Griffin¹

¹Francis Bitter National Magnet Laboratory and Department of Chemistry, Massachusetts Institute of Technology, Cambridge, Massachusetts 02139

²Plasma Fusion Center, Massachusetts Institute of Technology, Cambridge, Massachusetts 02139

³Department of Chemistry, Harvard University, Cambridge, Massachusetts 02138

(Received 26 July 1993)

DNP (dynamic nuclear polarization) experiments at 5 T are reported, in which a cyclotron resonance maser (gyrotron) is utilized as a 20 W, 140 GHz microwave source to perform the polarization. MAS (magic angle spinning) NMR spectroscopy with DNP has been performed on samples of polystyrene doped with the free radical BDPA (α, γ -bis(diphenylene- β -phenylallyl)) at room temperature. Maximal DNP enhancements of ~ 10 for ^1H and ~ 40 for ^{13}C are observed and are considerably larger than expected. The DNP and spin relaxation mechanisms that lead to these enhancements at 5 T are discussed.

1.5 % efficient

$(\gamma_e/\gamma_n) \sim 660$

$e^- \rightarrow ^1\text{H} \rightarrow ^{13}\text{C}$

$\epsilon \text{ max @ } \omega_e \pm \omega_n$

$e^- \rightarrow ^{13}\text{C}$

Courtesy, Griffin et al.

Direction des Sciences de la Matière

- High Frequency Dynamic Nuclear Polarization has been pioneered in the Griffin group @ MIT over the last two decades...

Massachusetts
Institute of
Technology

$T = 90 \text{ K}$
 $\nu_r = 8 \text{ kHz}$

DNP Enhanced ^{13}C MAS Spectra – ^{13}C Urea

Direction des Sciences de la Matière

Direction des Sciences de la Matière

Massachusetts
Institute of
Technology

Griffin group @ MIT

DNP up to 10 Tesla

 $\epsilon \sim 300 @ 90 \text{ K}$

Solid-state DNP: qualitative picture and challenges

Polarizing agents

- To increase $T_1\rho$, $T_s \sim 10\text{-}100\text{ K}$ are often required
- Use of cryoprotectant to obtain glassy matrix
 - avoid ice crystal formation
 - ensure uniform distribution of electrons
- DNP mechanism = e-H polarization transfer and relayed HH spin diffusion

Direction des Sciences de la Matière

Griffin's biradical

- EPR spectrum is 600 MHz wide @ 5 Tesla
Inhomogeneous broadening i.e. $\Delta \gg \omega_{0n}$
- Hole burning experiment on Tempo radicals
→ e-e couplings between the spin packets
- For CE: we need $\omega_{e1} - \omega_{e2} \sim \omega_n$
- Design of biradical in order to maximize the CE efficiency!
- For Totapol radical:
 - e-e coupling is 25 MHz
 - Electron concentration $\sim 10\text{ mM}$

K. Hu, Y.-y. Yu, T. Swager and R. G. Griffin, JACS, 2004, 126, 10844-10845

Direction des Sciences de la Matière

Polarization transfer to various systems

- Enhancement ~ 50 to 300 are routinely observed at 5 to 10 T and 90 K
- Development of optimized polarizing agents (e.g. TOTAPOL, BT2K)
- This approach has proven efficient in polarizing 100 nm size particles in 5 to 10 s

Direction des Sciences de la Matière

DNP in Amyloid Nanocrystals

Distribution of Polarization via ^1H Spin Diffusion

- When the dimensions of the system are large a polarization gradient develops !

Direction des Sciences de la Matière

Dissolution DNP

Direction des Sciences de la Matière

Dissolution DNP – ex situ DNP

Principle:

- DNP at moderate field and 1.4 K during ~1h
- dissolution step
- transfer to high field

The DNP ^{13}C polariser

3.35 T and 1.4K

Solid material doped with unpaired electrons

$P_e = 94\%$ and $P_c = 0.086\%$

J. Ardenkjær-Larsen, K. Golman et al., PNAS 100 (18), 10158 (2003).

Direction des Sciences de la Matière

The DNP ^{13}C polarizer

Challenges and limitations:

- Preserving the nuclear magnetization during the dissolution step
- Dilution factor
- Keep the magnetization during the transfer to high field magnet ($\rightarrow$ design of two centers magnet, etc.)
- one shot experiment

Main applications:

- ^{13}C hyperpolarization: small molecules structure determination, metabonomics, ...
- ^{13}C hyperpolarization for MRI: U- ^{13}C pyruvate etc...

Direction des Sciences de la Matière

Thank you for your attention

Direction des Sciences de la Matière