

Energy spectrum of primary cosmic rays in a wide range of energy

Janusz Kempa^{1,a}

¹Warsaw Univ. of Technology, Poland

Abstract. The works on this subject usually refer to the so-called. averaged spectra. The authors try to bypass the problem of inflating the spectra that result from other structure fluctuations which occur in each of the measuring device. The minimum particle spectrum of primary cosmic rays will be presented. The chemical composition of this radiation will be discussed.

See references [1–9]

References

- [1] E.G. Berezhko, *Advances in Space Research* 41 (2008) 429-441
- [2] E.S. Seo, V.S. Ptuskin, *Astrophysical Journal* 431 (1994) 705-714
- [3] V. Berezhinsky, A. Gazizov, S. Grigorieva, *Phys. Rev. D.* (2006) 74 043005
- [4] A. Garyaka et al, ISVHECRI Berlin 2012
- [5] V. Ptuskin, V. Zirakashvili, E.S. Seo, *Astrophysical Journal* 718 (2010) 31-36
- [6] E.G. Berezhko, S.P. Knurenko, L.T. Ksenofontov, *Astroparticale Physics* 36 (2010) 31-36
- [7] E.G. Berezhko, H. Voelk, *Astrophysical Journal* 661 (2007) L175-L178
- [8] Y.S. Yoon et al, arXiv:1102.2575v1 [astro-ph.HE] 13 Feb 2011
- [9] A.D. Panov et al, *Bulletin of the Russian Academy of Sciences Physics* 71 No 4 (2007) 494-497

Slides

The slides of the talk can be found on the website of the symposium ISVHECRI 2012:
<https://indico.desy.de/conferenceOtherViews.py?view=standard&confId=4594>

^akempa@pw.plock.pl