

Competition of β -delayed protons and β -delayed γ rays in ^{56}Zn and the exotic β -delayed γ -proton decay

S. E. A. ORRIGO^{1*}, B. RUBIO¹, Y. FUJITA², B. BLANK³,
W. GELLETLY⁴, J. AGRAMUNT¹, A. ALGORA^{1,5}, P. ASCHER³,
B. BILGIER⁶, L. CÁCERES⁷, R. B. ÇAKIRLI⁶, H. FUJITA⁸,
E. GANIOĞLU⁶, M. GERBAUX³, J. GIOVINAZZO³, S. GRÉVY³,
O. KAMALOU⁷, H. C. KOZER⁶, L. KUCUK⁶, T. KURTUKIAN-NIETO³,
F. MOLINA^{1,9}, L. POPESCU¹⁰, A. M. ROGERS¹¹, G. SUSOY⁶,
C. STODEL⁷, T. SUZUKI⁸, A. TAMII⁸ and J. C. THOMAS⁷

¹Instituto de Física Corpuscular, CSIC-Universidad de Valencia, E-46071
Valencia, Spain

²Department of Physics, Osaka University, Toyonaka, Osaka 560-0043, Japan

³Centre d'Etudes Nucléaires de Bordeaux Gradignan, CNRS/IN2P3 - Université
Bordeaux 1, 33175 Gradignan Cedex, France

⁴Department of Physics, University of Surrey, Guildford GU2 7XH, Surrey, UK

⁵Inst. of Nuclear Research of the Hung. Acad. of Sciences, Debrecen, H-4026,
Hungary

⁶Department of Physics, Istanbul University, Istanbul, 34134, Turkey

⁷Grand Accélérateur National d'Ions Lourds, BP 55027, F-14076 Caen, France

⁸Research Center for Nuclear Physics, Osaka University, Ibaraki, Osaka 567-0047,
Japan

⁹Comisión Chilena de Energía Nuclear, Casilla 188-D, Santiago, Chile

¹⁰SCK.CEN, Boeretang 200, 2400 Mol, Belgium

¹¹Physics Division, Argonne National Laboratory, Argonne, IL 60439, USA

Abstract

Remarkable results have been published recently on the β decay of ^{56}Zn . In particular, the rare and exotic β -delayed γ -proton emission has been detected for the first time in the fp shell. Here we focus the

*e-mail: sonja.orrigo@ific.uv.es

discussion on this exotic decay mode and on the observed competition between β -delayed protons and β -delayed γ rays from the Isobaric Analogue State.

1 Introduction

Decay spectroscopy is a powerful tool for exploring the structure of nuclei at the drip-lines. β -decay studies, in particular, provide direct access to the absolute values of the Fermi and Gamow-Teller transition strengths, $B(F)$ and $B(GT)$, respectively.

The proton-rich ^{56}Zn nucleus was observed for the first time at GANIL in 1999 [1]. ^{56}Zn is a weakly-bound nucleus lying very close to the proton drip-line. It has a quite small proton separation energy, $S_p = 560(140)$ keV [2], and third component of the isospin quantum number $T_z = -2$.

The first study of the β decay of ^{56}Zn was reported in *ref.* [3]. More recently, some interesting results on ^{56}Zn decay have been reported in *ref.* [4]. Among them the discovery of a rare and exotic decay mode, β -delayed γ -proton decay, which has been seen for the first time in the fp shell. The consequences of this rare decay sequence for the determination of the Gamow-Teller (GT) strength have also been analyzed.

2 The experiment

The experimental study of ^{56}Zn decay was performed at GANIL in 2010. The experiment used a primary beam of $^{58}\text{Ni}^{26+}$ to produce ^{56}Zn . The ^{58}Ni beam, of $3.7\text{ e}\mu\text{A}$ and accelerated to 74.5 MeV/nucleon , was fragmented on a natural Ni target, $200\text{ }\mu\text{m}$ thick. The fragments were selected by the LISE3 separator and implanted into a Double-Sided Silicon Strip Detector (DSSSD). The detection set-up comprised the aforementioned DSSSD detector, $300\text{ }\mu\text{m}$ thick, a silicon ΔE detector located 28 cm upstream, and four EXOGAM Ge clovers surrounding the DSSSD.

The EXOGAM clovers were used to detect β -delayed γ rays. The purpose of the DSSSD was the detection of both the implanted fragments and the subsequent charged-particle decays, *i.e.*, β particles and β -delayed protons. An implantation event was defined by simultaneous signals in both the ΔE and DSSSD detectors. A decay event was defined by a signal above threshold ($50\text{-}90\text{ keV}$) in the DSSSD and no coincident signal in the ΔE .

The implanted ions were identified and selected by putting a gate in a two-dimensional identification matrix, obtained by combining the energy

loss signal from the ΔE detector and the Time-of-Flight. The latter was defined as the time difference between the cyclotron radio-frequency and ΔE signal.

3 Results on the β decay of ^{56}Zn

The results on the β decay of ^{56}Zn [4] are summarized in the decay scheme in *fig. 1* and in table 1, and discussed below.

A half-life of $T_{1/2} = 32.9(8)$ ms was obtained for ^{56}Zn , in agreement with *ref.* [3]. To determine $T_{1/2}$, a decay-time spectrum has been constructed from the time correlations between a decay event in a given pixel of the DSSSD (with a total of 256 pixels) and any implantation signal that occurred before and after it in the same pixel, satisfying the identification condition required to select ^{56}Zn .




Figure 1: Scheme of the β decay of ^{56}Zn . The solid lines indicate observed proton or γ transitions, while the dashed lines correspond to transitions observed in the mirror ^{56}Co nucleus.

The analysis of the charged-particle spectrum measured in the DSSSD has provided new spectroscopic information on the energy levels populated in the ^{56}Cu nucleus, the β -daughter of ^{56}Zn . These levels are shown in *fig. 1*. The comparison of this level spectrum with that of the mirror ^{56}Co , obtained by the $^{56}\text{Fe}(^3\text{He},t)$ charge exchange reaction [5], has been very fruitful.

The analysis of the γ spectrum measured in the EXOGAM clovers and γ -proton coincidences have identified three γ rays at 309, 861 and 1835 keV.

Absolute $B(F)$ and $B(GT)$ strengths have been determined (table 1).

Table 1: β feedings, Fermi and Gamow Teller transition strengths to the ^{56}Cu levels populated in the β^+ decay of ^{56}Zn .

$E_X(\text{keV})$	$I_\beta(\%)$	$B(F)$	$B(GT)$
3508(140)*	43(5)	2.7(5)	
3423(140)	21(1)	1.3(5)	≤ 0.32
2661(140)	14(1)		0.34(6)
2537(140)	0		0
1691(140)	22(6)		0.30(9)
1391(140)	0		0

*Main component of the IAS.

3.1 Competition of β -delayed protons and β -delayed γ rays

In the first study of the ^{56}Zn β decay [3], the emission of β -delayed protons was observed but no β -delayed γ rays were seen. This was not a surprise because, in general, in proton-rich nuclei the proton decay is expected to dominate for states well above (>1 MeV) the proton separation energy S_p . The consequence is that normally the β feeding is directly inferred from the measured intensities of the proton peaks. However, cases where there is a competition between β -delayed proton emission and β -delayed γ de-excitation have also been observed, *e.g.*, in *refs.* [3, 6].

In the $T_z = -2 \rightarrow -1$, β^+ decay of ^{56}Zn to ^{56}Cu , the ^{56}Zn ground state decays with a Fermi transition to its Isobaric Analogue State (IAS) in ^{56}Cu . It should be noted that the de-excitation of this $T = 2$, $J^\pi = 0^+$ IAS via proton decay to the ground state of ^{55}Ni ($T = 1/2$, $J^\pi = 7/2^-$) is isospin forbidden. Therefore the proton emission that we observe can only happen through a $T = 1$ isospin impurity present in the IAS. Moreover in general, when the proton emission is isospin forbidden, the competitive emission of de-exciting γ rays from the IAS also becomes possible and can be observed even from IAS lying at an excitation energy well above S_p [3, 6].

The competition between β -delayed protons and γ rays has indeed been observed in ^{56}Zn . The γ decays represent 56(6)% of the total decays from the 3508 keV IAS. Thus one has to take into account the intensities of both the proton and γ peaks to determine the Fermi strength correctly.

We have also found evidence for the fragmentation of $B(F)$ due to a strong isospin mixing with a 0^+ state at 3423 keV [4], which is important in terms of the mass evaluation [7]. The isospin impurity in the ^{56}Cu IAS, $\alpha^2 = 33(10)\%$ (defined as in *ref.* [5]), and the off-diagonal matrix element of the charge-dependent part of the Hamiltonian, $\langle H_c \rangle = 40(23)$ keV, which is responsible for the isospin mixing of the 3508 keV IAS ($T = 2$, $J^\pi = 0^+$) and the 0^+ part of the 3423 keV level ($T = 1$), are similar to the values obtained in the mirror ^{56}Co nucleus [5].

Thus, the proton decay of the IAS proceeds thanks to the $T = 1$ component. However, considering the quite large isospin mixing in ^{56}Cu , the much faster proton decay ($t_{1/2} \sim 10^{-18}$ s) should dominate on the γ de-excitation ($t_{1/2} \sim 10^{-14}$ s in the mirror). This is not the case since we are still observing the γ decay of the IAS in competition with it.

The knowledge on the nuclear structure of the three nuclei involved in the decay, *i.e.*, ^{56}Zn , ^{56}Cu and ^{55}Ni , can provide us with a possible explanation for the hindrance of the proton decay. Shell model calculations are in progress to clarify this point.

3.2 The β -delayed γ -proton decay

Besides the competition between β -delayed proton emission and γ decay, the exotic sequence of β -delayed γ -proton decay has been detected. Indeed ^{56}Zn does β decay to its IAS in ^{56}Cu and from there we observe the emission of two γ rays of 861 and 1835 keV, populating the ^{56}Cu levels at 2661 and 1691 keV, respectively. Due to the low S_p , these levels are still proton-unbound and thereafter they decay by proton emission. Consequently the rare and exotic β -delayed γ -proton decay has been observed. In addition to these two branches, there is a third case. The 1691 keV level emits a γ ray of 309 keV, going to the level at 1391 keV that is again proton-unbound and then it de-excites by proton emission.

The β -delayed γ -proton decay has been observed here for the first time in the fp shell. This rare decay mode was seen only once before, in the sd shell in ^{32}Ar [6], but the consequences for the determination of $B(GT)$ were not addressed in *ref* [6].

The observation of this special decay mode is very important because it does affect the conventional way to determine $B(GT)$ near the proton drip-

line. For a proper determination of $B(\text{GT})$, indeed, it is crucial to correct the intensity of the proton transitions for the amount of indirect feeding coming from the γ de-excitation. This finding indicates that it is important to employ γ detectors in such studies. This decay mode is expected to be significant in heavier proton-rich nuclei with $T_z \leq -3/2$ under study at RIKEN.

Acknowledgements

This work was supported by the Spanish MICINN grants FPA2008-06419-C02-01, FPA2011-24553; Centro de Excelencia Severo Ochoa del IFIC SEV-2014-0398; CPAN Consolider-Ingenio 2010 Programme CSD2007-00042; *Junta para la Ampliación de Estudios* Programme (CSIC JAE-Doc contract) co-financed by FSE; MEXT, Japan 18540270 and 22540310; Japan-Spain coll. program of JSPS and CSIC; Istanbul University Scientific Research Projects, Num. 5808; UK Science and Technology Facilities Council (STFC) Grant No. ST/F012012/1; Region of Aquitaine. R.B.C. acknowledges support by the Alexander von Humboldt foundation and the Max-Planck-Partner Group. We acknowledge the EXOGAM collaboration for the use of their clover detectors.

References

- [1] J. Giovinazzo et al., *Eur. Phys. J. A*, **11** (2001) 247.
- [2] G. Audi et al., *Nucl. Phys. A*, **729** (2003) 1.
- [3] C. Dossat et al., *Nucl. Phys. A*, **792** (2007) 18.
- [4] S.E.A. Orrigo et al., *Phys. Rev. Lett.*, **112** (2014) 222501.
- [5] H. Fujita et al., *Phys. Rev. C*, **88** (2013) 054329.
- [6] M. Bhattacharya et al., *Phys. Rev. C*, **77** (2008) 065503.
- [7] M. MacCormick and G. Audi, *Nucl. Phys. A*, **925** (2014) 61.