The QCD equation of state at finite density from analytical continuation

Jana Günther^{1,a}, Rene Bellwied², Szabolcs Borsanyi¹, Zoltan Fodor^{1,3,4}, Sandor D. Katz^{3,5}, Attila Pasztor¹, and Claudia Ratti²

¹Department of Physics, University of Wuppertal, Gaussstr. 20, D-42119 Wuppertal, Germany

²Department of Physics, University of Houston, Houston, TX 77204, USA

³Inst. for Theoretical Physics, Eötvös University

⁴Jülich Supercomputing Centre, Forschungszentrum Jülich, D-52425 Jülich, Germany

⁵MTA-ELTE "Lendület" Lattice Gauge Theory Research Group, Budapest, Hungary

Abstract. An efficient way to study the QCD phase diagram at small finite density is to extrapolate thermodynamical observables from imaginary chemical potential. In this talk we present results on several observables for the equation of state to order $(\mu_B/T)^6$. The observables are calculated along the isentropic trajectories in the (T, μ_B) plane corresponding to the RHIC Beam Energy Scan collision energies. The simulations are performed at the physical mass for the light and strange quarks. μ_S was tuned in a way to enforce strangeness neutrality to match the experimental conditions; the results are continuum extrapolated using lattices of up to $N_t = 16$ temporal resolution.

1 Introduction

To analyse the quark gluon plasma that is created in heavy ion collision experiments at LHC or RHIC a theoretical understanding of the quark gluon plasma in QCD is needed. Around the deconfinement transition lattice simulations are the only systematic tool to study QCD since this region can not be accessed perturbatively. The investigation of the phase diagram at $\mu_B \neq 0$ is hindered by the sign problem. There have been several ideas in lattice QCD how to obtain results at real finite chemical potential like reweighting techniques [1–4], Taylor expansion [5–9], density of state methods [10, 11], using the canonical ensemble [12–14], the formulation with dual variables [15], Lefschetz thimbles [16, 17] or complex Langevin [18, 19]. Unfortunately these techniques are not yet able to solve the sign problem completely, therefore at the moment results that are continuum extrapolated and obtained at physical quark masses are restricted to vanishing or imaginary chemical potential. Nevertheless one needs information also at $\mu_B > 0$ particularly to analyse the collisions at RHIC [20]. In this work we use the method analytic continuation based on ensembles produced at imaginary chemical potential.

In terms of the baryon chemical potential μ_B the pressure P can be written as a function of temperature T :

$$\frac{P(\mu_B, T)}{T^4} = c_0(T) + c_2(T) \left(\frac{\mu_B}{T} \right)^2 + c_4(T) \left(\frac{\mu_B}{T} \right)^4 + c_6(T) \left(\frac{\mu_B}{T} \right)^6 + c_8(T) \left(\frac{\mu_B}{T} \right)^8 + O(\mu_B^{10}). \quad (1)$$

^ae-mail: Jana.Guenther@t-online.de

c_2 was calculated in the continuum in [21]. In [22] the coefficients c_2 to c_6 were presented on $N_t = 6$ and $N_t = 8$ lattices. In this proceedings we will go beyond those results and present the continuum extrapolated Taylor coefficients of the pressure up to c_6 based on [23]. Our analysis is done with

$$\langle n_S \rangle = 0 \text{ and } \langle n_Q \rangle = 0.4 \langle n_B \rangle. \quad (2)$$

2 Simulation details

We use a tree level Symanzik improved gauge action. The 2+1+1 dynamical flavours are simulated with four times stout smeared staggered fermions [24]. The smearing parameter is $\rho = 0.125$.

Our $T = 0$ simulations can be described as follows. The up and the down quarks have degenerate masses m_{ud} . Both m_{ud} and the strange quark mass m_s are tuned in a way that the pion and the kaon mass divided by the pion decay constant are set to their physical value for every lattice spacing. The charm mass m_c is set to be $m_c/m_s = 11.85$ as described in [25]. For setting the scale in physical units we use f_π or w_0 with lattice spacings between $a = 0.063$ fm and $a = 0.2$ fm. For all of our lattices $Lm_\pi > 4$ is fulfilled.

For the finite temperature runs we use non vanishing imaginary chemical potentials. Due to the Roberge-Weiss transition μ_B has to be smaller then $i\pi T$ in these runs (see for example [26–28]). Our values for μ_B are chosen as

$$\mu_B^{(j)} = iT \frac{j}{8} \pi \quad (3)$$

for $j \in \{0, 3, 4, 5, 6, 7\}$. We tune μ_S to maintain strangeness neutrality. More details on this procedure can be found in [29, 30]. For each value of μ_B we have simulations for sixteen temperatures between 135 MeV and 210 MeV on lattices with sizes $40^3 \times 10$, $48^3 \times 12$ and $64^3 \times 16$. In addition, we produced configurations with $j = 6.5$ on lattices with sizes $40^3 \times 10$ and $48^3 \times 12$. For each set of parameters we generated between 10000 and 20000 Hybrid Monte Carlo trajectories for the finite density configurations and 5 to 10 times more statistics for the $\mu_B = 0$ runs. For $N_t = 10$ and 12 every 5th configuration is included in our study. For $N_t = 16$ we analyse every 10th configuration. Though our simulations range up to a temperature of 210 MeV we use these ensembles to determine the equation of state by analytic continuation. In order to extend the equation of state up to approximately 300 MeV where the system is clearly in a state of quark gluon plasma, we collect data at these temperatures with $\mu_B = 0^1$. To reach 300 MeV we construct some *extrapolated data*. This means we are analysing a dataset

$$\left. \frac{d\hat{P}}{d\hat{\mu}_B} \right|_{\hat{\mu}_B} = \begin{cases} \left. \frac{d\hat{P}}{d\hat{\mu}_B} \right|_{\text{simulated at } \hat{\mu}_B} & \text{for } T < T_{\text{conn}} \\ \left. \frac{d\hat{P}}{d\hat{\mu}_B} \right|_{\text{extrapolated to } \hat{\mu}_B} & \text{for } T \geq T_{\text{conn}} \end{cases}. \quad (4)$$

Here the hat over a quantity means that it has been divided by the appropriate factors of T to make it dimensionless. To check whether it is safe to neglect higher order contributions we will use two different connection points T_{conn} from which on we will use the extrapolated data: One at 200 MeV and one at 220 MeV. The difference in this two analyses will then add to our systematic error.

¹In principle one may apply the Taylor expansion technique [21, 22] using these configurations. We did not follow this path.

3 Analysis

As the simulations on the different lattice sizes correspond to slightly different temperatures and we want to know the derivative with respect to T , a first step in the analysis is the interpolation of $\frac{n}{\mu_B T^2}$, $\frac{n_B}{\mu_B T^2}$ and $\frac{d\mu_Q}{d\mu_B}$ for every $\mu_B^{(j)}$. We use four different fit function:

$$A_1(T) = a + b(T - f) + \frac{c}{T - f} + d \arctan(e(T - f)) \quad (5)$$

$$A_2(T) = a + b(T - f) + \frac{c}{T - f} + \frac{d}{(1 + |e(T - f)|^g)^{1/g}}, \quad (6)$$

$$A_3(T) = a + b(T - f) + c(T - f)^2 + d \arctan(e(T - f)) \quad (7)$$

$$A_4(T) = a + b(T - f) + c(T - f)^2 + \frac{d}{(1 + |e(T - f)|^g)^{1/g}}. \quad (8)$$

The fit functions are purely meant as interpolation and can not be extended beyond the fit range. From here on we will analyse the different temperatures separately.

The next step in the analysis is to fit the data in the $\hat{\mu}_B^2$ direction. For this we use three different fit functions:

$$B_1(\hat{\mu}) = a + b\hat{\mu}^2 + c\hat{\mu}^4, \quad (9)$$

$$B_2(\hat{\mu}) = \frac{a + b\hat{\mu}^2}{1 + c\hat{\mu}^2}, \quad (10)$$

$$B_3(\hat{\mu}) = a + b\hat{\mu}^2 + c \frac{\sin(\hat{\mu})}{\hat{\mu}}. \quad (11)$$

For imaginary μ_B all three fit functions describe the data well. However, their extrapolation to real μ_B can be different as shown in figure 1. These differences are included in our systematic uncertainties. Having obtained the $\mu_B > 0$ data the next step is the continuum extrapolation. We use a linear fit as a function of $1/N_t^2$ though the data from $N_t = 10, 12$ and 16 . As an alternative analysis we also combine the fit in the μ_B direction and the continuum extrapolation. In this case we use the three fit functions

$$B_1(\hat{\mu}) = a_1 + \frac{a_2}{N_t^2} + \left(b_1 + \frac{b_2}{N_t^2}\right)\hat{\mu}^2 + \left(c_1 + \frac{c_2}{N_t^2}\right)\hat{\mu}^4, \quad (12)$$

$$B_2(\hat{\mu}) = \frac{a_1 + \frac{a_2}{N_t^2} + \left(b_1 + \frac{b_2}{N_t^2}\right)\hat{\mu}^2}{1 + \left(c_1 + \frac{c_2}{N_t^2}\right)\hat{\mu}^2}, \quad (13)$$

$$B_3(\hat{\mu}) = a_1 + \frac{a_2}{N_t^2} + \left(b_1 + \frac{b_2}{N_t^2}\right)\hat{\mu}^2 + \left(c_1 + \frac{c_2}{N_t^2}\right) \frac{\sin(\hat{\mu})}{\hat{\mu}}. \quad (14)$$

From equation (9) to (11) or from equation (12) to (14) we can determine the Taylor coefficients of the pressure. The results are shown in figure 2. The statistical error is determined via the bootstrap method with 1000 bootstrap samples.

Figure 1. The increasing error at larger μ_B from the three different fit functions depending on the temperature. While at $\mu_B < 0$ the fit functions look very similar they can still differ at positive μ_B . Here the data from the $N_t = 12$ lattice is shown as this allows a comparison to the raw data.

For the systematic error we use the histogram method of [31, 32]. In this case we have the following options to chose from for different analyses:

- two methods of scale setting
- four fits in the T direction
- three fits in the μ_B direction
- combined and separate μ_B -fit and continuum extrapolation
- two different points T_{conn} from which we use the data extrapolated from $\mu_B = 0$ at high temperatures.

This adds up to a total of $2 \cdot 4 \cdot 3 \cdot 2 \cdot 2 = 96$ analyses. The systematic uncertainty is defined by the central 68% of the distribution. Figure 2 shows the result with statistical and the systematic error added quadratically.

The importance of the higher order terms can be seen from figure 3. From the Taylor coefficient we can calculate $\hat{n}_B$ to different orders. Though c_2 gives the pressure up to $O(\mu_B^2)$ it provides $\hat{n}_B$ only up to $O(\mu_B)$. The reason for that is that $\hat{n}_B$ is obtained as a derivative with respect to μ_B . For a temperature of 145 MeV only the c_2 and c_4 coefficient contribute to the result because c_6 is tiny. At this temperature the analytic continuation for $\hat{n}_B$ up to order $O(\mu_B^5)$ only increases the error compared to the previous order. This is not surprising as figure 2 shows that at $T = 145$ MeV c_6 is compatible with zero. However at a temperature where c_6 differs significantly from zero, for example 170 MeV, the additional order has a visible influence.

Besides the Taylor coefficients $c_2 \dots c_6$ we compute the equation of state. In heavy ion collisions if one neglects dissipative effects, the quark gluon plasma created in the collision expands with a fixed

Figure 2. The Taylor coefficients of the pressure. The values for c_0 are taken from [33]. The red line shows the expectations of the HRG model.

baryon number and without generation of entropy. Therefore the ratio $\frac{S}{N_B}$ is constant. The medium cools down along these isentropic trajectories in the T - μ_B -plane. Therefore we compute the equation of state along these trajectories. In figure 4 we show six of them which were matched to the beam energies of RHIC as determined in [34]. The solid lines are predictions of the HRG model that we use to continue our trajectories at lower temperatures. The $\frac{S}{N_B} = 30$ choice corresponds to a beam energy of 14.5 GeV. As it can be seen from the errors this is at the end of how far we can extrapolate in μ_B . Since the error for this case is relatively large we compute the equation of state along the trajectories with $\frac{S}{N_B} = 51$ and $\frac{S}{N_B} = 420$. Figure 5 shows the pressure and the trace anomaly along these isentropic trajectories.

4 Conclusion

We investigated the equation of state around the QCD transition temperature at finite baryon chemical potential with lattice QCD. Since this can not be done directly due to the sign problem, we used simulations at imaginary baryon chemical potential to analytically continue to real chemical potentials. We determined the Taylor coefficients of the pressure up to $\mathcal{O}(\mu_B^6)$ in the continuum limit. Results up to $\mu_B/T \approx 2$ were given. All simulations were done at the strangeness neutral point to match conditions in heavy ion collision experiments. We then calculated the pressure and the trace anomaly along the isentropic trajectories. The trajectories were matched to the RHIC beam energies.

Figure 3. The contribution of the different order coefficients to the baryon number. At 145 MeV the contribution of c_6 is negligible while at 170 MeV it influences the results.

Figure 4. The trajectories of constant $\frac{S}{N_B}$ matched to the beam energies of RHIC as determined in [34].

Figure 5. The pressure and the trace anomaly along trajectories of constant $\frac{S}{N}$.

Acknowledgements

C.R. would like to thank Volker Koch, Jacquelyn Noronha-Hostler, Jorge Noronha and Bjorn Schenke for fruitful discussions. This project was funded by the DFG grant SFB/TR55. This material is based upon work supported by the National Science Foundation through grant number NSF PHY-1513864 and by the U.S. Department of Energy, Office of Science, Office of Nuclear Physics, within the framework of the Beam Energy Scan Theory (BEST) Topical Collaboration. An award of computer time was provided by the INCITE program. This research used resources of the Argonne Leadership Computing Facility, which is a DOE Office of Science User Facility supported under Contract DE-AC02-06CH11357. The authors gratefully acknowledge the Gauss Centre for Supercomputing (GCS) for providing computing time for a GCS Large-Scale Project on the GCS share of the supercomputer JUQUEEN [35] at Jülich Supercomputing Centre (JSC).

References

- [1] I.M. Barbour, S.E. Morrison, E.G. Klepfish, J.B. Kogut, M.P. Lombardo, Nucl. Phys. Proc. Suppl. **60A**, 220 (1998), hep-lat/9705042
- [2] Z. Fodor, S.D. Katz, Phys. Lett. **B534**, 87 (2002), hep-lat/0104001
- [3] Z. Fodor, S.D. Katz, JHEP **03**, 014 (2002), hep-lat/0106002
- [4] F. Csikor, G.I. Egri, Z. Fodor, S.D. Katz, K.K. Szabo, A.I. Toth, Nucl. Phys. Proc. Suppl. **119**, 547 (2003), [547(2002)], hep-lat/0209114
- [5] C.R. Allton, S. Ejiri, S.J. Hands, O. Kaczmarek, F. Karsch, E. Laermann, C. Schmidt, L. Scorzato, Phys. Rev. **D66**, 074507 (2002), hep-lat/0204010
- [6] C.R. Allton, M. Doring, S. Ejiri, S.J. Hands, O. Kaczmarek, F. Karsch, E. Laermann, K. Redlich, Phys. Rev. **D71**, 054508 (2005), hep-lat/0501030
- [7] R.V. Gavai, S. Gupta, Phys. Rev. **D78**, 114503 (2008), 0806.2233
- [8] S. Basak et al. (MILC), PoS **LATTICE2008**, 171 (2008), 0910.0276
- [9] O. Kaczmarek, F. Karsch, E. Laermann, C. Miao, S. Mukherjee, P. Petreczky, C. Schmidt, W. Soeldner, W. Unger, Phys. Rev. **D83**, 014504 (2011), 1011.3130
- [10] Z. Fodor, S.D. Katz, C. Schmidt, JHEP **03**, 121 (2007), hep-lat/0701022
- [11] A. Alexandru, C. Gattringer, H.P. Schadler, K. Splittorff, J.J.M. Verbaarschot, Phys. Rev. **D91**, 074501 (2015), 1411.4143
- [12] A. Alexandru, M. Faber, I. Horvath, K.F. Liu, Phys. Rev. **D72**, 114513 (2005), hep-lat/0507020
- [13] S. Kratochvila, P. de Forcrand, PoS **LAT2005**, 167 (2006), hep-lat/0509143
- [14] S. Ejiri, Phys. Rev. **D78**, 074507 (2008), 0804.3227
- [15] C. Gattringer, PoS **LATTICE2013**, 002 (2014), 1401.7788
- [16] L. Scorzato, PoS **LATTICE2015**, 016 (2016), 1512.08039
- [17] A. Alexandru, G. Basar, P. Bedaque, Phys. Rev. **D93**, 014504 (2016), 1510.03258
- [18] E. Seiler, D. Sexty, I.O. Stamatescu, Phys. Lett. **B723**, 213 (2013), 1211.3709
- [19] D. Sexty, Phys. Lett. **B729**, 108 (2014), 1307.7748
- [20] G. Odyniec, EPJ Web Conf. **95**, 03027 (2015)
- [21] S. Borsanyi, G. Endrodi, Z. Fodor, S.D. Katz, S. Krieg, C. Ratti, K.K. Szabo, JHEP **08**, 053 (2012), 1204.6710
- [22] P. Hegde (BNL–Bielefeld–CCNU), Nucl. Phys. **A931**, 851 (2014), 1408.6305

- [23] J. Gunther, R. Bellwied, S. Borsanyi, Z. Fodor, S.D. Katz, A. Pasztor, C. Ratti (2016), **1607.02493**
- [24] C. Morningstar, M.J. Peardon, Phys. Rev. **D69**, 054501 (2004), hep-lat/**0311018**
- [25] C. McNeile, C.T.H. Davies, E. Follana, K. Hornbostel, G.P. Lepage, Phys. Rev. **D82**, 034512 (2010), **1004.4285**
- [26] M. D’Elia, F. Sanfilippo, Phys. Rev. **D80**, 111501 (2009), **0909.0254**
- [27] C. Pinke, O. Philipsen, PoS **LATTICE2013**, 209 (2014)
- [28] L.K. Wu, X.F. Meng, Phys. Rev. **D90**, 094506 (2014), **1405.2425**
- [29] R. Bellwied, S. Borsanyi, Z. Fodor, J. Günther, S.D. Katz, C. Ratti, K.K. Szabo, Phys. Lett. **B751**, 559 (2015), **1507.07510**
- [30] J. Günther, R. Bellwied, S. Borsanyi, Z. Fodor, S.D. Katz, C. Ratti, K. Szabo, PoS **LATTICE2015**, 142 (2016)
- [31] S. Durr et al., Science **322**, 1224 (2008), **0906.3599**
- [32] S. Borsanyi et al., Science **347**, 1452 (2015), **1406.4088**
- [33] S. Borsanyi, Z. Fodor, C. Hoelbling, S.D. Katz, S. Krieg, K.K. Szabo, Phys. Lett. **B730**, 99 (2014), **1309.5258**
- [34] P. Alba, W. Alberico, R. Bellwied, M. Bluhm, V. Mantovani Sarti, M. Nahrgang, C. Ratti, Phys. Lett. **B738**, 305 (2014), **1403.4903**
- [35] Jülich Supercomputing Centre, Journal of large-scale research facilities **A1**, 1 (2015)