Numerical study of the electron and muon lateral distribution in atmospheric showers of high energy cosmic rays

Atreidis Georgios^{1,a}

¹ *Physics Department, Aristotle University of Thessaloniki, 54124 Thessaloniki, Greece, - Nuclear and Elementary Particle Physics Division*

Abstract. The lateral distribution of an atmospheric shower depends on the characteristics of the high energy interactions and the type of the primary particle. The influence of the primary particle in the secondary development of the shower into the atmosphere, is studied by analyzing the lateral distribution of electron and muon showers having as primary particle, proton, photon or iron nucleus. This study of the lateral distribution can provide useful conclusions for the mass and energy of the primary particle. This paper compares the data that we get from simulations with CORSIKA program with experimental data and the theoretical NKG function expressing lateral electron and muon distribution. Then we modify the original NKG function to fit better to the simulation data and propose a method for determining the mass of the original particle started the atmospheric shower.

1. The simulation

The Monte Carlo technique is used for the simulation of the lateral growth of an extensive air shower which has a specified primary particle type and energy corresponding to a specific interaction model. The CORSIKA (COsmic Ray Simulations for KAscade) is a Monte Carlo simulation program for the physics of cosmic rays, used by many cosmic ray experiments for various actions. It can be used to simulate interactions and decays of nuclei, hadrons, muons, electrons, and photons in the atmosphere up to energies of some 10^{20} eV. It gives details for all secondary particles that are created in an air shower and pass a selected observation level.

In this paper, for each entry angle we ran 100 simulations with fixed primary particle energy and of zenith angles varying from 0 to 75 degrees. The results obtained as the mean value of all simulations for each particle. The corresponding errors are due to the limited number of events in the Monte Carlo simulation.

^ae-mail: gatrdis@physics.auth.gr

Table 1. Simulation data

magnitude	value
Shower sets	10
Shower/set	100
Primary particle	Proton, Helium, Gamma, Iron
Primary energy	10 ¹³ eV – 10 ¹⁶ eV
Zenith angle	0° to 75°
Azimuth angle	-180° to 180°
Energy cut (hadrons, muons)	0.3 GeV
Energy cut (electrons, photons)	0.003 GeV
Starting height	0 g/cm ²
Earth’s magnetic field Thessaloniki	B _x =25.005μT, B _z =39.488μT
Observation level	110 m

2. Lateral Distribution of Electrons

Compare of the lateral distribution of electrons for different primary particle and at different angles.

When the entry angle θ is small $0^\circ < \theta < 20^\circ$ the differences of lateral distribution (of electrons arriving at the observation plane and the electrons arriving at the plane perpendicular to the axis of the cascade) are small. But when the entry angle increases, some electrons arrive at the observation level (Earth's surface) by the shortest path, while others arrive from the longest path.

The cascade development on the short path is faster. In the long path we have greater absorption of electrons, thus the larger the entry angle; the fewer electrons reach at this point.

The lateral distribution of electrons for primary particle proton, photon and iron nuclei, for 0 degrees entry angle is shown in Figure 1. Respective Figures derived (according to the simulation) for small entry angles $0^\circ < \theta < 20^\circ$.

Figure 1. Lateral distribution of electrons for cascades of primary particles proton, photon and iron nuclei with primary energy 10⁷ GeV and entry angle 0 degrees.

But in all the scenarios of the simulation where the primary particle is photon (electromagnetic cascade), the density of electrons is greater. The diagram of Figure 1 shows that there is symmetry around the center of the cascade (in a plane perpendicular to the axis of the cascade).When the entry

angle is small $\theta < 20$ degrees, the plane which is perpendicular to the axis of the cascade and the observation plane are almost identical. But as the entry angle increases, the angle of the two planes increases respectively. Thus there is no longer symmetry on both sides of the axis of the cascade at the level of observation.

The variation in the number of lateral electron explained by the difference of the two roads (for angle $\theta = 60$ degrees), followed by the electrons in the cascade. Thus, the lateral distribution of electrons is not symmetrical (Figure 2).

What causes this?

When increasing the entry angle, the cascade develops into greater heights, since we have more lateral route to the higher layers of the atmosphere. So the electrons arriving at observation level are significantly less. But over 40 degrees and apparently over 60 degrees the data differ in the symmetry of the lateral distribution. The electrons that reach the observation level from the long path are less than those arriving from the small path (Figure 2).

In very large zenith angles $\theta \sim 90$ degrees, cosmic rays develop their own cascades in the upper atmosphere, with a similar mechanism as that in the vertical cascades. Their electromagnetic component almost completely absorbed by the increased atmospheric side depth and cannot reach the ground. A few electrons with irregular distribution reach the Earth's surface from only one side of the cascade. In contrast, the high energy neutrinos can cause the creation of horizontal cascades at a larger depth in the atmosphere and therefore likened to vertical atmospheric cascades as to the particles that reach the ground.

So the assumption that we can detect high energy neutrinos by horizontal cascades takes on a consistent basis.

Figure 2. In large entry angles $\theta > 60^\circ$ for electrons arriving at observation level there is no symmetry. The cascade develops differently in the lateral plane.

Compare of the lateral distribution of electrons on the same original particle and at different angles.

We applied the simulation for a proton primary particle with energy 10^7 GeV and for different entry angles. By increasing the entry angle, the number of electrons that reach the ground decreased, accordingly to the reasons mentioned above.

Another reason for the reduction of the lateral electron is the height of the first interaction. Obviously in large entry angles that height increases since the initial particle travels deep side to first interaction.

Figure 3. Lateral distribution of electrons for cascades with a proton primary particle with primary energy 10^7 GeV and entry angles 0, 20 and 40 degrees.

3. Lateral Distribution of muons

The number of muons arriving at the observation level is growing by the energy of the primary particle. Figure 4 shows the comparison of the lateral distribution of electron and muon at an atmospheric shower with primary particle a proton with energy 10^7 GeV. In the same picture is also shown the lateral distribution of muons from the data of the KASCADE experiment. As it is known, the lateral distribution of muons at the sea level and in the region of energy considered in this work is much more flat and typically one order of magnitude lower than the lateral distribution of electrons in our own scale distance from the core.

Figure 4. Lateral electron and muon distribution for atmospheric shower with a primary particle proton with energy 10^7 GeV. - Lateral muon distribution for atmospheric showers with a primary particle proton, helium and iron nucleus. The primary energy is 10^7 GeV.

4. The NKG function

The NKG (Nishimura-Kamata-Greisen) function is a good theoretical approach to the lateral distribution of electrons and muons.

The NKG function for electrons

The function which is adapted to the electrons, is given by the formula below.

$$\rho(r) = N_e C(s) \left(\frac{r}{r_M}\right)^{s-2} \left(1 + \frac{r}{r_M}\right)^{s-4.5} \qquad C(s) = \frac{\Gamma(4.5-s)}{2\pi r_M^2 \Gamma(s) \Gamma(4.5-2s)}$$

Where C(s) is normalization constant, r_M is the Moliere radius (for electrons the Moliere radius is 80m in the atmosphere) and s is the time slope parameter (age parameter).

However, based on knowledge of previews studies, it seems that the NKG functions have shortcomings in the implementation of the measured lateral distribution of electrons, becoming more evident in the large distances from the axis of the cascade. This deficit is usually caused by the fact that the NKG function has occurred for electromagnetic cascades while hadronic cascades are a superposition of a large number of independent electromagnetic cascades. For typical cascade, detectors extend in approximately 200m from the axis of the cascade.

The NKG function for muons

The function which is adapted to the muons, is given by the formula below.

$$\rho(r) = N_e C(s) \left(\frac{r}{r_M}\right)^{s-2} \left(1 + \frac{r}{r_M}\right)^{s-4} \qquad C(s) = \frac{\Gamma(4-s)}{2\pi r_M^2 \Gamma(s) \Gamma(4-2s)}$$

Where C(s) is normalization constant, r_M is the Moliere radius (for muons the Moliere radius is 420m in the atmosphere) and s is the time slope parameter (age parameter).

5. Reconstruction of Cosmic Rays for the Determination of their Mass

The results obtained from the simulation of atmospheric cascades with CORSIKA lead us to the modification of the NKG function by varying the exponent values. The new function which fits better the results of the simulation than the original NKG function is:

For electrons

$$\rho = C \left(\frac{r}{r_o}\right)^{s-1.8} \left(1 + \frac{r}{r_o}\right)^{2s-3.8} \qquad \text{And} \qquad C = \frac{N_e}{2\pi r_o^2} \frac{\Gamma(3.8-s)}{\Gamma(s+0.2) \Gamma(3.6-2s)}$$

For muons

$$\rho = C \left(\frac{r}{r_o}\right)^{s-1.6} \left(1 + \frac{r}{r_o}\right)^{s-3.8} \qquad \text{And} \qquad C = \frac{N_e}{2\pi r_o^2} \frac{\Gamma(3.8-s)}{\Gamma(s+0.4) \Gamma(3.4-2s)}$$

Table 2. Exponent s for electrons

Particle - Nucleus	Exponent s
Proton	0.75±0.06
Helium	0.63±0.06
Oxygen	0.79±0.06
Silicon	0.85±0.06
Calcium	0.89±0.06
Iron	0.90±0.06

Table 3. Exponent s for muons

Particle - Nucleus	Exponent s
Proton	0.724±0.062
Helium	0.616±0.061
Oxygen	0.757±0.063
Silicon	0.828±0.061
Calcium	0.874±0.065
Iron	0.896±0.067

The table 2 and 3 give the exponent s obtained as the mean value of all simulations for each particle. The corresponding errors are due to the limited number of events in the Monte Carlo simulation.

Figure 5. Lateral distribution of electrons for cascades with a primary particle proton having primary energy 10^7 GeV and entry angle 20 degrees. The red line represents the fitted function while the blue represents the original. - Muon lateral distribution for atmospheric showers with primary particle helium nucleus. The blue line is the NKG function. The red line is the modified NKG function.

Figure 6. Variation of the exponent s (electrons – muons), in relation to the mass of the primary particle.

References

[1] ELSEVIER, W.D. Apel et al., “Comparison of measured and simulated lateral distributions for electrons and muons with KASCADE,” *Astroparticle Physics* 24, 467–483, 2006

[2] KASCADE Collaboration, T. Antoni et al., “Electron, muon and hadron lateral distributions measured in air showers by the KASCADE experiment,” *Astroparticle Physics* 14, 245-260, 2001

[3] 32nd Int. Cosmic Ray Conf., D. Kalita, K.Boruah, “Study of Lateral distribution Parameters from simulation of HE Cosmic Ray EAS,” Vol. 1, 247-250, 2011

[4] D. Heck - T. Pierog, “Extensive Air Shower Simulation with CORSIKA, A User’s Guide”, 2013