

Search for time-reversal-invariance violation in double polarized antiproton-deuteron scattering

Yuriy Uzikov^{1,2,3,*} and Johann Haidenbauer^{4,**}

¹Joint Institute for Nuclear Research, Dubna 141980, Russia

²Dubna State University, Dubna 141980, Russia

³Department of Physics, M.V. Lomonosov Moscow State University, Moscow 119991, Russia

⁴Institute for Advanced Simulation and Institut für Kernphysik, Forschungszentrum Jülich GmbH, D-52428 Jülich, Germany

Abstract.

Apart from the pd reaction also the scattering of antiprotons with transversal polarization p_y^p on deuterons with tensor polarization P_{xz} provides a null-test signal for time-reversal-invariance violating but parity conserving effects. Assuming that the time-reversal-invariance violating $\bar{N}N$ interaction contains the same operator structure as the NN interaction, we discuss the energy dependence of the null-test signal in $\bar{p}d$ scattering on the basis of a calculation within the spin-dependent Glauber theory at beam energies of 50-300 MeV.

1 Introduction

Under CPT symmetry time-reversal-invariance violating but parity conserving (TVPC) forces are considered as a possible source of CP-invariance violation, which is required to account for the matter-antimatter asymmetry in the universe [1]. In contrast to effects from time-reversal-invariance violation together with parity violation such as a permanent electric dipole moment (EDM) of elementary particles, so far much less attention was paid to TVPC effects. The reason why TVPC effects are interesting is that experimental limits on them are still rather weak, in particular, considerably weaker than those for the EDM.

Since the intensity of TVPC interactions within the standard model is extremely small [2], an observation of any effects at the present accuracy level of experiments would be a direct indication of physics beyond the standard model. Indeed a pertinent measurement is planned at the COSY accelerator in the Research Center in Jülich [3]. The observable in question is the integrated cross section for scattering of protons with transversal polarization p_y^p on deuterons with tensor polarization P_{xz} . It provides a null-test signal for TVPC effects [4] and it will be measured in pd scattering at 135 MeV [3]. Theoretical studies of the energy dependence of the expected signal were performed at energies of the planned experiment [5–12] on the basis of the spin-dependent Glauber theory and demonstrate several unexpected effects. Among them are (i) the absence of the contribution from the

*e-mail: uzikov@jinr.ru

**e-mail: j.haidenbauer@fz-juelich.de

lowest-mass meson-exchange (ρ meson) in the TVPC NN interaction, caused by its specific isospin, spin and momentum dependence; (ii) a strong impact of the deuteron D -wave on the null-test signal due to a destructive interference between the S - and D -wave contributions, even for zero transferred 3-momentum; (iii) oscillating behaviour of the null-test signal as a function of the beam energy, i.e. the vanishing of the TVPC signal at some specific energies is possible even when the TVPC interaction itself is nonzero; (iv) a very small influence of the Coulomb interaction on the TVPC term of the pd forward scattering amplitude \bar{g} . Furthermore, certain relations between differential observables of elastic pd scattering caused by time-reversal-invariance requirements were obtained and the degree of their violation by TVPC NN forces was studied [13, 14].

Since the spin structure of the amplitude for pd - and $\bar{p}d$ elastic scattering is the same, it is obvious that the integrated cross section for scattering of a polarized ($p_y^{\bar{p}}$) antiproton on tensor polarized (P_{xz}) deuterons also provides a null-test signal for TVPC effects. Furthermore, the TVPC $\bar{N}N$ amplitude for elastic scattering contains the same operator structures as the one for TVPC NN elastic scattering, except for the charge-exchange terms. Therefore, the formalism developed in Refs. [7, 8, 11] within the Glauber theory for the calculation of the null-test signal in pd scattering can be straightforwardly applied to $\bar{p}d$ scattering too. However, due to differences in the hadronic part of the pN and $\bar{p}N$ scattering amplitudes and also in the electromagnetic interactions, the energy dependence of the null test signal in pd and $\bar{p}d$ interaction has to be different. In the present work the energy dependence of the null-test signal in $\bar{p}d$ scattering is studied on the basis of calculations within the spin-dependent Glauber theory using the spin-dependent $\bar{p}N$ amplitudes from a recent partial wave analysis of $\bar{p}p$ scattering [15].

2 Null-test signal for time-reversal-invariance violation

The total cross section for $\bar{p}d$ scattering with TVPC forces included can be written in the same form as for pd scattering [7]

$$\sigma_{tot} = \sigma'_0 + \sigma'_1 \mathbf{p}^{\bar{p}} \cdot \mathbf{p}^d + \sigma'_2 (\mathbf{p}^{\bar{p}} \cdot \mathbf{m})(\mathbf{p}^d \cdot \mathbf{m}) + \sigma'_3 P_{zz} + \bar{\sigma} p_y^{\bar{p}} P_{xz}^d. \quad (1)$$

Here $\mathbf{p}^{\bar{p}}$ (\mathbf{p}^d) is the vector polarization of the initial antiproton (deuteron), P_{zz} and P_{xz} are the tensor polarizations of the deuteron, and $p_y^{\bar{p}}$ is the transversal component of the antiproton vector polarization. The OZ axis is directed along the beam direction \mathbf{m} , the OY axis is directed along the vector polarization of the antiproton beam $\mathbf{p}^{\bar{p}}$ and the OX axis is chosen to form a right-handed reference frame. The integrated cross sections σ'_i ($i = 0, 1, 2, 3$) are those which arise from a standard time-reversal invariant and parity conserving interaction, while the last term $\bar{\sigma}$ appears only in the presence of the TVPC interactions and constitutes the TVPC null-test signal. The result (1) can be derived using phenomenological $\bar{p}d$ forward scattering amplitudes and the generalized optical theorem.

The evaluation of the integrated cross sections σ'_i and $\bar{\sigma}$ at beam energies > 100 MeV can be done on the basis of the spin-dependent Glauber theory of $\bar{p}d$ scattering which is formulated similarly to the theory of pd scattering given in Ref. [16]. Indeed, as shown in Ref. [17], this theory allows one to describe rather well available data on differential spin observables of pd scattering in the forward hemisphere at beam energies of 135 – 200 MeV. For the antiproton-deuteron scattering this theory can be applied at even lower energies due to the presence of strong annihilation effects. In the Glauber theory one uses the elastic (on-shell) $\bar{N}N$ scattering amplitudes as input. Hadronic amplitudes of the $\bar{p}N$ scattering are taken here in the same form as for pN scattering [16]

$$M_N(\mathbf{p}, \mathbf{q}; \sigma, \sigma_N) = A_N + C_N \sigma \hat{\mathbf{n}} + C'_N \sigma_N \hat{\mathbf{n}} + B_N(\sigma \hat{\mathbf{k}})(\sigma_N \hat{\mathbf{k}}) + (G_N + H_N)(\sigma \hat{\mathbf{q}})(\sigma_N \hat{\mathbf{q}}) + (G_N - H_N)(\sigma \hat{\mathbf{n}})(\sigma_N \hat{\mathbf{n}}), \quad (2)$$

where $\hat{\mathbf{q}}$, $\hat{\mathbf{k}}$ and $\hat{\mathbf{n}}$ are defined as unit vectors along the vectors $\mathbf{q} = (\mathbf{p} - \mathbf{p}')$, $\mathbf{k} = (\mathbf{p} + \mathbf{p}')$ and $\mathbf{n} = [\mathbf{k} \times \mathbf{q}]$, respectively; \mathbf{p} (\mathbf{p}') is the initial (final) antiproton momentum.

In general, the TVPC NN interaction contains 18 different terms [18]. In the case of the on-shell NN scattering amplitude there are only three terms with different (independent) spin-momentum structures. In the present study we consider the following two terms for the TVPC (on-shell) t -matrix of elastic $\bar{p}N$ scattering which have the same structure as those in TVPC pN scattering

$$t_{\bar{p}N} = h_N[(\boldsymbol{\sigma} \cdot \mathbf{k})(\boldsymbol{\sigma}_N \cdot \mathbf{q}) + (\boldsymbol{\sigma}_N \cdot \mathbf{k})(\boldsymbol{\sigma} \cdot \mathbf{q}) - \frac{2}{3}(\boldsymbol{\sigma}_N \cdot \boldsymbol{\sigma})(\mathbf{k} \cdot \mathbf{q})]/m_p^2 + \quad (3)$$

$$+ g_N[\boldsymbol{\sigma} \times \boldsymbol{\sigma}_N] \cdot [\mathbf{q} \times \mathbf{k}][\boldsymbol{\tau} - \boldsymbol{\tau}_N]_z/m_p^2.$$

Here $\boldsymbol{\sigma}$ ($\boldsymbol{\sigma}_N$) is the Pauli matrix acting on the spin state of the antiproton (nucleon $N = p, n$) and $\boldsymbol{\tau}$ ($\boldsymbol{\tau}_N$) is the isospin matrix acting on the isospin state of the antiproton (nucleon). The momenta \mathbf{q} and \mathbf{k} were already defined above in the context of Eq. (2). Both terms in Eq. (3), h_N and g_N , occur in the TVPC pn interaction. The TVPC pN scattering amplitude contains also the charge-exchange term

$$t^{ch} = g'_N(\boldsymbol{\sigma} - \boldsymbol{\sigma}_N) \cdot i[\mathbf{q} \times \mathbf{k}][\boldsymbol{\tau} \times \boldsymbol{\tau}_N]_z/m_p^2, \quad (4)$$

which describes the elastic transitions $pn \rightarrow np$ and $np \rightarrow pn$. Within a picture of one-meson-exchange interaction this g' -term corresponds to the charged ρ -meson exchange [19]. The same term (4) corresponds to the charge-exchange processes $\bar{p}p \rightarrow \bar{n}n$ or $\bar{n}n \rightarrow \bar{p}p$. However, in contrast to pn scattering these processes are inelastic and therefore the operation of time-reversal invariance transforms, for example, the $\bar{p}p \rightarrow \bar{n}n$ amplitude to the $\bar{n}n \rightarrow \bar{p}p$ amplitude and does not impose any restrictions on these amplitudes.

The h_N -term in Eq. (3) can be associated with the axial h_1 -meson exchange. As shown in Ref. [19], contributions of the π - and σ -meson to the TVPC NN interaction are excluded, which is obviously true for the TVPC $\bar{N}N$ interaction as well.

2.1 TVPC amplitude of $\bar{p}d$ forward scattering

One can write the $\bar{p}d$ forward elastic scattering amplitude in general form taking into account the TVPC $\bar{N}N$ interactions, as it was done for pd elastic scattering [7, 17], and then apply the generalized optical theorem to derive Eq. (1) for the total $\bar{p}d$ scattering cross section. As in Ref. [7], the integrated cross section $\bar{\sigma}$ is related to the TVPC term \bar{g} of the $\bar{p}d$ forward elastic scattering amplitude by $\bar{\sigma} = -4\sqrt{\pi} \text{Im} \frac{2}{3} \bar{g}$. Furthermore, the TVPC forward amplitude of $\bar{p}d$ elastic scattering \bar{g} can be found within the Glauber theory [7]. We consider the h_N - and g_N -terms and take into account both the S - and D -wave components of the deuteron. Taking into account that the g_N -term is excluded in the process $\bar{p}n \rightarrow \bar{p}n$ due to the isospin operator in Eq. (3), we obtain the following result for the TVPC forward amplitude from the corresponding equation in Ref. [11]:

$$\bar{g} = \frac{i}{4\pi m_p} \int_0^\infty dq q^2 \left[S_0^{(0)}(q) - \sqrt{8} S_2^{(1)}(q) - 4 S_0^{(2)}(q) + \sqrt{2} \frac{4}{3} S_2^{(2)}(q) + 9 S_1^{(2)}(q) \right] \quad (5)$$

$$[-C'_n(q)(h_p + g_p) - C'_p(q)h_n].$$

Here $S_i^{(j)}$ are the elastic form factors of the deuteron defined in Ref. [11]. The first term in the (big) squared brackets in Eq. (5), $S_0^{(0)}(q)$, corresponds to the S -wave approximation, the second term, $S_2^{(1)}(q)$, accounts for the S - D interference, and the last three terms contain the pure D -wave contributions.




Figure 1. The TVPC signal $\tilde{\sigma}$ for the h -term, in units of the ratio (ϕ_h , see Ref. [7]) of the TVPC and the strong $h_1 NN$ coupling constants, versus the antiproton beam energy T . Results of our calculations accounting for different terms of the deuteron wave function in Eq. (5) are shown, based on the deuteron wave function of the CD Bonn potential and the hadronic $\bar{p}N$ amplitudes from Ref. [15]: S -wave (black), S - D interference (blue), $S + S$ - D waves (green), full result (red).

As was shown in Ref. [11], the contribution of the g' -term to the null-test signal vanishes in pd scattering due to the specific spin-isospin structure of the g' -interaction. Formally, for the same reason the charge-exchange g' -term given by Eq. (4) vanishes in the $\bar{p}d$ forward elastic scattering amplitude.

In the first theoretical work [5] where the null-test signal was calculated within the impulse approximation, the Coulomb interaction was not considered. In Ref. [6] Faddeev calculations were performed, but only for nd scattering and at rather low energies of ~ 100 keV. The Coulomb interaction was taken into account for the first time in Ref. [7] in a calculation of the null-test signal of pd scattering within Glauber theory and found to be negligible. A similar result was found in Ref. [20] using Faddeev calculations.

2.2 Numerical results

Results of numerical calculations of the energy dependence of the null test-signal for the h -term are presented in Fig. 1, in units of the unknown TVPC coupling strength. One can see from this figure that the deuteron S -wave contribution (dashed line) leads to a smooth energy dependence and has a node at an antiproton beam energy of about 50 MeV. The inclusion of the D -wave changes this behaviour considerably (solid line) due to a destructive S - D interference (cf. dash-dotted line). As a result, a second zero of the null-test signal $\tilde{\sigma}$ appears at higher energies, i.e. at $T \approx 300$ MeV. The maximal value of $\tilde{\sigma}$ is expected at 100 – 150 MeV. Note that the actual position of the nodes changes only slightly when deuteron wave functions from other NN models are used for the calculation.

Let us consider possible spurious effects that could mimic a TVPC signal. One source for a spurious signal is associated with a nonzero deuteron vector polarization $p_d^y \neq 0$ (in the direction of the incident-proton-beam polarization \mathbf{p}^p). In this case, the term $\sigma_1 P_y^{\bar{p}} p_y^d$ in Eq. (1) contributes to the

asymmetry corresponding to the difference of the event counting rates for the cases of $p_y^{\bar{p}} P_{xz} > 0$ and $p_y^{\bar{p}} P_{xz} < 0$ (with the fixed sign of P_{xz}), which is planned to be measured at COSY [3]. According to our calculations, the integrated cross section σ_1 could be equal to zero at antiproton beam energies of ~ 100 MeV (see results for the Jülich $\bar{N}N$ interaction model in Refs. [21, 22]). Therefore, at this energy the spurious signal caused by a nonzero value of the deuteron vector polarization p_y^d could be minimized.

3 Concluding remarks

We have performed a study of time-reversal-invariance violating but parity conserving effects in antiproton-deuteron scattering. Specifically, we have evaluated the null-test TVPC signal for scattering of antiprotons with transversal polarization $p_y^{\bar{p}}$ on deuterons with tensor polarization P_{xz} on the basis of the spin-dependent Glauber theory. The observed effects turned out to be similar to those in pd scattering: (i) There is a strong impact of the deuteron D -wave on the null-test signal that arises from a destructive interference between the S - and D -wave contributions; (ii) There is an oscillating behaviour of the null-test signal as a function of the beam energy. Accordingly, it is possible that the signal for TVPC effects is zero at some specific energies, even when the TVPC interaction itself is nonzero.

Acknowledgement. This work was supported in part by the Heisenberg-Landau program.

References

- [1] A.D. Sakharov, Pis'ma v ZHETF **5**, 32 (1967).
- [2] R.S. Conti, I.B. Khriplovich, Phys. Rev. Lett. **68**, 3262 (1992).
- [3] COSY Proposal N 215, Spokespersons: P.D. Eversheim, B. Lorentz, and Yu. Valdau, http://apps.fz-juelich.de/pax/paxwiki/images/8/8c/215-TRI_Prop_sum.pdf
- [4] H.E. Conzett, Phys. Rev. C **48**, 423 (1993).
- [5] M. Beyer, Nucl. Phys. A **560**, 895 (1993).
- [6] Y.-H. Song, R. Lazauskas, V. Gudkov, Phys. Rev. C **84**, 025501 (2011).
- [7] Yu.N. Uzikov, A.A. Temerbayev, Phys. Rev. C **92**, 014002 (2015).
- [8] Yu. N. Uzikov, J. Phys. Conf. Ser. **678**, 012020 (2016).
- [9] Yu.N. Uzikov, EPJ Web Conf. **113**, 04027 (2016).
- [10] Yu.N. Uzikov, A.A. Temerbayev, Int. J. Mod. Phys. Conf. Ser. **40**, 1660080 (2016).
- [11] Yu.N. Uzikov, J. Haidenbauer, Phys. Rev. C **94**, 035501 (2016).
- [12] Yu.N. Uzikov, EPJ Web Conf. **138**, 08001 (2017).
- [13] A.A. Temerbayev, Yu. N. Uzikov, Izv. Ross. Akad. Nauk Ser. Fiz. **79**, 535 (2015) [Bull. Russ. Acad. Sci. Phys. **79**, 493 (2015)].
- [14] A.A. Temerbayev, Yu. N. Uzikov, Izv. Ross. Akad. Nauk Ser. Fiz. **80**, 271 (2016) [Bull. Russ. Acad. Sci. Phys. **80** N4, 242 (2016)].
- [15] D. Zhou, R.G.E. Timmermans, Phys. Rev. C **86**, 044003 (2012).
- [16] M.N. Platonova, V.I. Kukulín, Phys. Rev. C **81**, 014004 (2010).
- [17] A.A. Temerbayev, Yu.N. Uzikov, Yad. Fiz. **78**, 38 (2015) [Phys. At. Nucl. **78**, 35 (2015)].
- [18] P. Herczeg, Nucl. Phys. **75**, 655 (1966).
- [19] M. Simonius, Phys. Lett. **B 78**, 147 (1975).
- [20] Y.-H. Song, R. Lazauskas, V. Gudkov, Phys. Rev. C **93**, 065501 (2016).

- [21] Yu. N. Uzikov, J. Haidenbauer, Phys. Rev. C **87**, 054003 (2013).
- [22] Yu. N. Uzikov, J. Haidenbauer, Phys. Rev. C **88**, 027001 (2013).