

Resonances in positron scattering on a supercritical nucleus and spontaneous production of e^+e^- pairs

Sergey Godunov^{1,*} Bruno Machet^{2,**} and Mikhail Vysotsky^{1,***}

¹ITEP

²UPMC (LPTHE)

Abstract. Solving the Dirac equation for a positron in the Coulomb field of a nucleus with $Z > Z_{\text{cr}}$ we observe (confirm) the resonance dependence of the scattering phase δ_z on the energy ε_p of the positron. A physical interpretation of these resonances is suggested.

1 Introduction

This talk is based on [1].

The Dirac equation in the Coulomb field of a nucleus for the ground energy level gives

$$E_0 = m \cdot \sqrt{1 - (Z\alpha)^2}.$$

What happens when $Z > 1/\alpha = 137$?

When nucleus radius is accounted for, the square root singularity is resolved, with growing Z ground state energy becomes negative [2].

At $Z = Z_{\text{cr}} \approx 175$ ground state energy dives into lower continuum of the Dirac equation, E_0 becomes less than $-m$.

As soon as a nucleus with $Z > Z_{\text{cr}}$ is formed, spontaneous production of e^+e^- pairs occurs; positrons are radiated to spatial infinity and can be detected while electrons are captured by the nucleus. Naked supercritical nucleus is screened by electrons [3], [4], [5], [6], [7].

2 Several physical phenomena

QCD

Gribov theory of quark confinement: growing of the strong (color) charge at large distances leads to quark pair creation from vacuum. Color charge is screened and color objects do not exist as asymptotic states (confinement).

Gravity

Strong gravitational potential leads to particle creation - black hole radiates.

No experimental confirmation of the QED phenomenon up to present time!
(Some hints from graphene physics.)

*e-mail: sgodunov@itep.ru

**e-mail: machet@lpthe.jussieu.fr

***e-mail: vysotsky@itep.ru

3 The recent development

The paper by V.M. Kuleshov et al. was published recently [8].

Main points of [8]:

1. Resonances in positron scattering on supercritical nucleus
WE CONFIRM IT
2. Stability of the naked nucleus with arbitrary large Z ; no e^+e^- pairs creation
WE DO NOT AGREE

4 The Rutherford scattering

1, 2, 3, 4 volumes of Landau and Lifshitz (from Mechanics to QED) contain considerations of the Rutherford scattering, however for $Z > 137$ new phenomena occurs.

The energy levels in the Dirac equation becomes complex.

These resonances were briefly discussed in [6], [9], [10], [11], [12].




Figure 1. Complex ϵ plane

5 The Dirac equation

The radial functions of the Dirac equation $F(r) \equiv rf(r)$ and $G(r) \equiv rg(r)$ are determined by the following differential equations:

$$\begin{cases} \frac{dF}{dr} + \frac{\kappa}{r}F - (\epsilon + m - V(r))G = 0, \\ \frac{dG}{dr} - \frac{\kappa}{r}G + (\epsilon - m - V(r))F = 0. \end{cases}$$

According to the paper by I. Pomeranchuk and Ya. Smorodinsky, to deal with $Z\alpha > 1$ case we choose:

$$V(r) = -\frac{Z\alpha}{r}, \quad r > R,$$

$$-\frac{Z\alpha}{R}, \quad r < R.$$

Solution of the Dirac equation.

Solution at $r < R$:

$$\begin{pmatrix} F \\ G \end{pmatrix} = \text{const} \cdot \sqrt{\beta r} \cdot \begin{pmatrix} \mp J_{\mp(1/2+\kappa)}(\beta r) \\ J_{\pm(1/2-\kappa)}(\beta r) \frac{\beta}{\varepsilon+m+\frac{Z\alpha}{R}} \end{pmatrix},$$

where $\kappa = \pm 1, \pm 2, \dots, \beta = \sqrt{(\varepsilon + \frac{Z\alpha}{R})^2 - m^2}$. Upper (lower) signs correspond to $\kappa \leq 0$.

Solution at $r > R$:

$$\begin{pmatrix} F \\ G \end{pmatrix} = \begin{pmatrix} \sqrt{m+\varepsilon} \\ -\sqrt{m-\varepsilon} \end{pmatrix} \exp(-\rho/2) \rho^{i\tau} \begin{pmatrix} Q_1 + Q_2 \\ Q_1 - Q_2 \end{pmatrix},$$

where $\tau = \sqrt{(Z\alpha)^2 - \kappa^2}, \rho = 2\lambda r = -2ikr, \lambda = \sqrt{(m-\varepsilon)(m+\varepsilon)}$, and

$$\left\{ \begin{array}{l} Q_1 = C \cdot \frac{-\frac{iZ\alpha m}{k} + \kappa}{-i\tau + \frac{iZ\alpha\varepsilon}{k}} \cdot {}_1F_1\left(i\tau - \frac{iZ\alpha\varepsilon}{k}, 2i\tau + 1, \rho\right) + \\ \quad + D \cdot \frac{-\frac{iZ\alpha m}{k} + \kappa}{i\tau + \frac{iZ\alpha\varepsilon}{k}} \rho^{-2i\tau} {}_1F_1\left(-i\tau - \frac{iZ\alpha\varepsilon}{k}, -2i\tau + 1, \rho\right), \\ Q_2 = C \cdot {}_1F_1\left(1 + i\tau - \frac{iZ\alpha\varepsilon}{k}, 2i\tau + 1, \rho\right) + D \cdot \rho^{-2i\tau} {}_1F_1\left(1 - i\tau - \frac{iZ\alpha\varepsilon}{k}, -2i\tau + 1, \rho\right), \end{array} \right.$$

where C and D are arbitrary coefficients. Unlike textbooks case both regular and singular at $r = 0$ solutions are taken into account.

The ratio C/D should be determined by matching with the solution at $r < R$.

6 Scattering phase and the position of the poles

The scattering phase $\delta_\kappa(\varepsilon, Z)$ is determined by investigating the behavior of the wave function at large r :

$$\begin{pmatrix} F \\ G \end{pmatrix} \Big|_{r \rightarrow \infty} \propto \begin{pmatrix} \sqrt{m+\varepsilon} \\ -\sqrt{m-\varepsilon} \end{pmatrix} \left\{ e^{i(kr + \frac{Z\alpha\varepsilon}{k} \ln(2kr))} e^{2i\delta} \pm e^{-i(kr + \frac{Z\alpha\varepsilon}{k} \ln(2kr))} \right\},$$

$$e^{2i\delta_\kappa} = -\frac{1}{\kappa + \frac{iZ\alpha m}{k}} \cdot \frac{\frac{C}{D} \cdot \frac{\Gamma(2i\tau)}{\Gamma(i\tau + \frac{iZ\alpha\varepsilon}{k})} \rho^{i\tau} (-\rho)^{-i\tau} - \frac{\Gamma(-2i\tau)}{\Gamma(-i\tau + \frac{iZ\alpha\varepsilon}{k})} \rho^{-i\tau} (-\rho)^{i\tau}}{\frac{C}{D} \cdot \frac{\Gamma(2i\tau)}{\Gamma(1+i\tau - \frac{iZ\alpha\varepsilon}{k})} - \frac{\Gamma(-2i\tau)}{\Gamma(1-i\tau - \frac{iZ\alpha\varepsilon}{k})}}.$$

The resonance of the scattering amplitude corresponds to the pole of the S -matrix element $S \equiv e^{2i\delta}$, so we immediately get an equation for the position of this pole in the ε -plane:

$$\frac{C}{D} \cdot \frac{\Gamma(2i\tau)}{\Gamma(1+i\tau - \frac{iZ\alpha\varepsilon}{k})} - \frac{\Gamma(-2i\tau)}{\Gamma(1-i\tau - \frac{iZ\alpha\varepsilon}{k})} = 0.$$




Figure 2. Dependence on ε_p of the scattering phase $\delta_{-1}(\varepsilon_p, 232)$ ($Z = 232$ and $\kappa = -1$) for a nucleus with radius $R = 0.031/m$. The blue solid line corresponds to the exact phase, the green dashed line corresponds to the approximation $1/m \gg R$ in matching, a’la *Kuleshov et al.*

Where do resonances (Gamov states) come from?

- At $Z < Z_{cr}$ (ground state energy $\varepsilon > -m$) they are bound states, spectrum of which are given by zeroes of denominator of S .
- At $Z > Z_{cr}$ pole goes to the second sheet of energy plane ABOVE left cut,

$$\varepsilon = -\varepsilon_0 + \frac{i}{2}\gamma$$

Kuleshov et al.:

At $Z > Z_{cr}$ state is at lower continuum – sign of energy should be reversed (negative electron energies at lower continuum describe positive positron energies).

BUT: resonance state originates from upper continuum ...

7 The Dirac equation for positron

The Dirac equation for positron in the field of (positively charged) nucleus

$$Z\alpha \longrightarrow -Z\alpha, \quad \tilde{V}(r) = -V(r)$$

Solving this equation we found scattering phase δ .

What about resonances?




Figure 3. The dependence of the ground state energy on Z . The square markers are for the exact values of the energy and the round markers are for the approximate ones. At $Z = Z_{cr}$ the bound states become resonances with positive $\text{Im}[\varepsilon]$.

At first glance – no bound states, so no resonances.

However we should not forget, that the Dirac equation describes BOTH electrons and positrons. Thus “the Dirac equation for positron” describes BOTH positrons and electrons.

It describes bound states, which this time originates from lower continuum, where electrons are situated.

At small Z the ground state energy equals

$$\varepsilon = -m(1 - (Z\alpha)^2/2).$$

With growing Z energy crosses zero and at $Z > Z_{cr}$ first resonance (Gamov) state emerges. It is situated on the second sheet of energy plane below right cut:

$$\varepsilon_{ps} = \varepsilon_0 - \frac{i}{2}\gamma.$$

It is a proper place for the resonance state!

Qualitative explanation of the resonance phenomena in the e^+N^+ system; M.I. Eides question.

A Schrödinger-like equation from the Dirac equation for positron:

$$\chi'' + k^2\chi = 0,$$

where $k^2 = 2m(E - U)$, $E = \frac{\varepsilon^2 - m^2}{2m}$. The effective potential is seen to be made of two terms: $U = U_1 + U_2$, where:

$$U_1 = \frac{\varepsilon}{m} \tilde{V} - \frac{1}{2m} \tilde{V}^2 - \frac{\varkappa(1 - \varkappa)}{2mr^2},$$

and

$$U_2 = \frac{\tilde{V}''}{4m(\varepsilon - m - \tilde{V})} + \frac{3}{8m} \frac{(\tilde{V}')^2}{(\varepsilon - m - \tilde{V})^2} + \frac{\varkappa \tilde{V}'}{2mr(\varepsilon - m - \tilde{V})}.$$

for $\varepsilon = m$ and $\varkappa = 1$, we get

$$U = \frac{Z\alpha}{r} - \frac{4(Z\alpha)^2 - 3}{8mr^2}.$$

At short distances the terms $\propto 1/r^2$ dominates and, for a supercritical nucleus, they lead to attraction, while the Coulomb term dominates at $r \geq 1/m$. This attractive force explains the existence of resonances in the e^+N^+ system. For $Z\alpha > 1$ fall to the center phenomena occurs.

8 Conclusions

According to V.M. Kuleshov et al. since the scattering phase is real e^+e^- pairs are not produced. That's why naked nucleus with arbitrary large Z is stable.

OUR OPINION: No naked nucleus with $Z > Z_{cr}$.

- Conservation of energy allows pairs production.
- Resonance in the heavy nucleus - positron system just signal pair production.
- What happens:
 - Without nucleus empty level in lower continuum is positron.
 - With nucleus the energy of empty level gets imaginary part, so it decays with the lifetime $1/\gamma$.
 - In this time electron which occupies the state with the same energy in the lower continuum “jumps” to this empty level creating “charged vacuum”, while positron (hole in the lower continuum) is radiated to “infinity”.




Figure 4. Electron-positron pair production.

1. Rutherford scattering of positron on supercritical nucleus do exhibit resonance behavior; width of this resonance equals time in which e^+e^- pair is produced by supercritical nucleus.

2. *Kuleshov et al.* wrote in Conclusions of their paper, that stable state of the supercritical nucleus consists of empty states of upper continuum, empty discrete levels and occupied states of lower continuum. It is just the picture I described. Occupied by electrons states of lower continuum which come from the upper continuum when Z grows and becomes larger than Z_{cr} form “charged vacuum” – its charge equals $-n$, where n is the number of these levels. Compensating charge $+n$ was carried out by produced positrons.
3. Conclusive evidence: Let us consider positron scattering on a nucleus with, say, $Z = Z_{cr} + 10$. Two electrons from charged vacuum are spread at $r \approx 1/m$. Resonance in this case will be approximately at the same place which we found before having approximately the same width. This width gives the lifetime of positron in vicinity of supercritical nucleus; during this time positron flies away leaving nuclei screened by vacuum electrons.

Supercritical nucleus is not naked - its electric charge is partially screened by these electrons.

We are very grateful to the organizers of the QUARKS-2018 for the great event.
S.G. and M.V. are supported by RFBR under grant 16-02-00342.

References

- [1] S.I. Godunov, B. Machet and M.I. Vysotsky, *Eur. Phys. J. C* **77**, 782 (2017).
- [2] I. Pomeranchuk and Ya. Smorodinsky, *J. Phys. USSR* **9**, 97 (1945).
- [3] V.V. Voronkov and N.N. Kolesnikov, *JETP* **12**, 136 (1961).
- [4] S.S. Gershtein and Ya.B. Zel'dovich, *JETP* **30**, 358 (1970).
- [5] W. Pieper and W. Greiner, *Z. Phys.* **218**, 327 (1969).
- [6] Ya.B. Zeldovich and V.S. Popov, *UFN* **105**, 403 (1971).
- [7] W. Greiner, B. Müller, and J. Rafelski, *Quantum Electrodynamics of Strong Fields* (Springer, Berlin, 1985).
- [8] V.M. Kuleshov, V.D. Mur, N.B. Narozhny, A.M. Fedotov, Yu.E. Lozovik, and V.S. Popov, *Phys. Usp.* **58**, 785 (2015).
- [9] B. Müller, J. Rafelski, and W. Greiner, *Z. Phys.* **257**, 62 (1972).
- [10] B. Müller, J. Rafelski, and W. Greiner, *Z. Phys.* **257**, 183 (1972).
- [11] V.D. Mur and V.S. Popov, *Theor. Math. Phys.* **27**, 429 (1976).
- [12] V.S. Popov, V.L. Eletsky, and V.D. Mur, *Sov. Phys. JETP* **44**, 451 (1976).