


# High-power Microwaves Against Locusts and Other Harmful Animals

V.E.Zapevalov

Institute of Applied Physics RAS, Nizhny Novgorod, Russia, zapev@appl.sci-nnov.ru

Locusts, acridia - some species of insects of the present locust family (Acrididae), capable of forming large flocks (up to hundreds of millions of individuals) migrating to considerable distances [1-3]. The damage it causes to plants, incomparable. Locusts can destroy all the plants in its path, and its invasion has always been a universal scourge. Innumerable flocks of these insects destroyed the crops, which inevitably led to mass starvation. A feature of biology of the locust is the presence of two phases - single and herd. Herd and single locust phases have significant differences, both in appearance and physiology, and in the nature of behavior. Locusts of a single phase usually have a protective coloration, a well-defined sexual dimorphism, and a low-active solitary lifestyle. Insects of the gregarious phase are colored more brightly and contrastively, adults are more adapted to flight. Locusts in this phase behave much more actively are formed larvae or adults. There is practically no sexual dimorphism in the herd phase. Locust has in its life cycle such stages: eggs, then larvae, the latter – imago (fig.1). The larvae moult 5 times and after the final moult they have wings, the imago stage begins.

The hordes of locusts are able to move at a speed of 15-20 km/h and fly without interruption for up to 20 hours in a row. Huge living clouds of locusts sometimes reach 10 km in width and up to 200 km in length. Two to three days of tailwind is enough for locusts to attack even very remote areas.



**Fig. 1.** Locusts: egg capsules; larvae of the 1st- 5th instar; adult insect.


The fight against this evil is conducted worldwide, but almost all used methods have a strong negative impact on the environment [1-3]. Of course, in the late XX -early XXI centuries researchers have the opportunity to obtain fundamentally new data on these insects with the help of molecular-genetic, biochemical and information methods. This especially applies to the mechanisms of the transition from a single phase to a herd and back, migrations of swarms and flocks, and so on. However, these opportunities are often not implemented. This is largely due to the

fact that interest in these insects (as well as research funding) falls sharply after the suppression of another outbreak, when the danger for agriculture passes. The ultimate goal in locust control is the use of preventive and proactive methods that disrupt the environment to the least possible extent. This would make agricultural production easier and more secure in the many regions where growing crops is of vital importance to the survival of the local people.

In this publication we are going to demonstrate that for control and suppression of locust (and other harmful animals) it is appropriate to use of high-power microwave systems. Important issues are the selection of the frequency range and the required power level for creating mobile microwave complexes. These conditions are determined by technical and economic factors, availability and cost of necessary sources of microwave radiation, transportation (radiation and propagation) of electromagnetic waves, detection of insects and influence on egg capsules, kuligas (larvae), flocks (imago).


The characteristic number of individuals in the migratory flock of locusts is  $N \sim 10^7$ - $10^9$  pieces with a density of individuals  $n \sim 10^3$ - $10^4$  m<sup>-3</sup>. Mass of locust specimen is near  $\sim 3$  g. The characteristic value of biomass in the flock is up to 3000 tons. The time of flock in flight is about  $\sim 12$  hours per day. Locust locating can be remotely conducted by infrared radiation using thermal imagers since the temperature of the clusters is distinctly higher than the underlying surface. Radar and visual methods can be used also.

Important issue is the proper selection of the frequency range and required power levels for exposure to different insects. In the microwave range, the cell membranes become practically short-circuited. Therefore, the electrical properties of suspensions of cells and tissues with high water content are determined by the aqueous, salt and protein content of the intra- and intercellular environment. Fig.2 shows dielectric properties of water which determine absorption of microwaves (pure water and with salt) as function of wavelength [4, 5].



**Fig. 2** Absorption of microwaves by water (pure - solid lines, and with salt - dashed lines).


Strong absorption of microwaves by water at the body of insects (pure and mainly with salt as hemolymph) effects on egg-caps, larvae and flocks of imago. In view of the specific structure of insects (fig. 3), effects of microwaves on them are much stronger than for mammals. All vital locust organs are located close to the surface [1-3], within the skin layer, in contrast to mammals (and in general vertebrates) where 83% of the energy of this radiation is absorbed by the upper layer of the skin.



**Fig. 3.** Scheme of locust and its main organs

It is advisable for effects on eggs and larvae to use a high-efficient magnetrons and gyrotrons [6, 7] at the ISM frequencies (0.915, 2.45, 5.8, 24.125 GHz etc.). Requirements for such type devices for complexes against locust could be discussed. It is reasonable that due to technical and economical reason preferable power level now is 1-3 kW for magnetron (total power of complex near 100 kW) and order of 100 kW for gyrotron.

In adult stage (imago) locust clusters the clear advantages have gyrotron systems of mm range similar non-lethal weapons [8]. Fig. 4 shows vehicle-mounted Active Denial System (ADS). This advanced system uses CPI's VGB-8095 gyrotron to generate a 95 GHz, 100 kW beam which provides a safe but effective deterrent. Similar Chinese system, known as the Poly WB-1, uses millimeter-wave beams (near 30 GHz) to scald targets from up to a kilometer away.




**Fig. 4.** ADS and gyrotron with cryomagnet.

It could be promising to use gyrotron operating at the harmonics of cyclotron frequency [7-10]. Fig. 5 demonstrates power and efficiency of gyrotron at the 2<sup>nd</sup> harmonics of cyclotron frequency (the  $TE_{03}$ ,  $n=2$ ,  $\lambda_0 \approx 12$  mm,  $f_0 \approx 25$  GHz) vs. beam current. Some other acceptable version of gyrotrons for such system described at [10]. For those systems RF beam control by phased array antenna it will be practically useful pos-

sibility of frequency tuning of RF source [7-8].

There are a lot of other phytophagous and harmful animals of agriculture and forestry in our country and the entire world. The possibility of using similar microwave systems to combat other harmful animals, pests of agriculture and forestry could be developed also.



**Fig. 5.** Power and efficiency of 2<sup>nd</sup> harmonics gyrotron.

### Summary

It is reasonable to use high-power microwaves to control locusts and other harmful animals.

Locust locating can be remotely conducted using thermal imagers because the temperature of the clusters is noticeably higher than the underlying surface. Due to the peculiarities of the insects structure the microwaves influence for them is much stronger than for mammals.

To effect on the laying of eggs and larvae, it is advisable to use high-efficiency magnetrons and gyrotrons at ISM frequencies. At the adult stage, gyrotron systems, analogous to non-lethal weapons, have clear advantages. The most attractive versions of magnetron and gyrotrons complexes for above mentioned tasks were pointed out.

### References

- [1] Resh V.H. (ed.), Carde R.T. (ed.). Encyclopedia of Insects. Academic Press. 2003. p.666-669.
- [2] A. Steedman, (ed). 1990. Locust Handbook (3rd edition). Chatham: Natural Resources Institute. 204 pp.
- [3] B.Uvarov, 1966. Grasshoppers and locusts, V. I. Cambridge: University Press. 481 pp.
- [4] J. B. Hasted, Liquid water: Dielectric properties, in Water. A comprehensive treatise, Vol 1, Ed. F. Franks (Plenum Press, New York, 1972) pp. 255-309.
- [5] Sh.E. Tsimring, Electron beams and microwave vacuum electronics, John Wiley and Sons, Inc., Hoboken, New Jersey, 2007.
- [6] M.Thumm. State-of-the-Art of High Power Gyro-Devices and Free Electron Masers. KIT Scientific Publishing, 2016.
- [7] V.E.Zapevalov, Evolution of gyrotrons, Radiophysics and Quantum electronics, 2011, v. LIV, №8-9, pp. 559-572.
- [8] V.E.Zapevalov, S.A.Malygin, Sh.E.Tsimring, Gyrotrons at the second harmonic of the cyclotron frequency. In the book: Gyrotrons, Gorky. IAP, 1980, pp.171-187.
- [9] A.N.Kuftin, V.A.Flyagin, V.K.Lygin, et al. 5.8-62 GHz cw gyrotrons with warm and permanent magnets for technological application. Proc. of Int. workshop "Strong Microwaves in Plasmas", 1999, N. Novgorod, 2000. pp. 671-676.