

# Comment on the $X(3915)$ nonstandard hadron candidate

Stephen Lars Olsen<sup>1,\*</sup>

<sup>1</sup>University of Chinese Academy of Science, Beijing 100049, CHINA

**Abstract.** I review the experimental evidence for the  $X(3915)$ , the candidate nonstandard meson associated with  $\omega J/\psi$  resonance-like peaks in  $B \rightarrow K\omega J/\psi$  and  $\gamma\gamma \rightarrow \omega J/\psi$  near  $M(\omega J/\psi) = 3920$  MeV, and address the conjecture that it can be identified as the  $\chi'_{c2}$ , the radial excitation of the  $\chi_{c2}$  charmonium state. Since the partial decay width for  $B \rightarrow KX(3915)$  is at least an order-of-magnitude higher than that for  $B \rightarrow K\chi_{c2}$ , its assignment as the  $\chi'_{c2}$  is dubious.

## 1 Introduction

A number of meson candidates, dubbed the XYZ mesons, that contain charmed-quark anticharmed-quark ( $c\bar{c}$ ) pairs but do not match expectations for any of the unassigned levels of the  $[c\bar{c}]$  charmonium meson spectrum, have been observed in recent experiments [1]. In some cases, the distinction between the new states that are nonstandard hadrons and conventional charmonium mesons remains controversial.

This is especially the case for the  $X(3915)$  that was first observed by Belle [2] and confirmed by BaBar [3, 4] as a near-threshold peak in the  $\omega J/\psi$  invariant mass distribution in exclusive  $B \rightarrow K\omega J/\psi$  decays (see Fig. 1a). An  $\omega J/\psi$  mass peak with similar mass and width was seen in the two-photon fusion process  $\gamma\gamma \rightarrow \omega J/\psi$ , again by both Belle [5] and BaBar [6] (see Fig. 1b); BaBar reported its  $J^{PC}$  to be  $0^{++}$ . The similar masses and widths of the peaks seen in the two production modes suggest that these are being produced a single state (i.e., the  $X(3915)$ ). The Particle Data Group's (PDG) average values for the mass and width measurements from both production channels are [7]:

$$M(X(3915)) = 3918.4 \pm 1.9 \text{ MeV} \quad \text{and} \quad \Gamma(X(3915)) = 20.0 \pm 5.0 \text{ MeV}, \quad (1)$$

and the product branching fraction for  $X(3915)$  production in  $B^+$  meson decays is


$$\mathcal{B}(B^+ \rightarrow K^+ X(3915)) \times \mathcal{B}(X \rightarrow \omega J/\psi) = 3.0 \pm 0.9 \times 10^{-5}. \quad (2)$$

The measured  $\gamma\gamma \rightarrow \omega J/\psi$  production rates are used to extract the ( $J^{PC}$ -dependent) widths:

$$\Gamma_{\gamma\gamma}(X(3915)) \times \mathcal{B}(X \rightarrow \omega J/\psi) = 54 \pm 9 \text{ eV} (0^{++}) \quad \text{or} \quad 11.4 \pm 2.7 \text{ eV} (2^{++}). \quad (3)$$

---

\*e-mail: salsensnu@gmail.com




**Figure 1.** a) The  $\omega J/\psi$  invariant mass spectrum for  $B \rightarrow K\omega J/\psi$  decays from (top) Belle [2] and (bottom) Babar [4]. The low mass peak in the BaBar data is attributed to  $X(3872) \rightarrow \omega J/\psi$  (see inset); the higher mass peak is the  $X(3915) \rightarrow \omega J/\psi$  signal. The Belle analysis did not consider the possible presence of an  $X(3872) \rightarrow \omega J/\psi$  signal. b) The  $\omega J/\psi$  mass spectrum for  $\gamma\gamma \rightarrow \omega J/\psi$  from (top) Belle [5] and (bottom) Babar [6].

## 2 The $X(3915)$ is not the $\chi'_{c0}$ charmonium state?

The Babar group's  $J^{PC}$  determination was based on an analysis of angular correlations amongst the final-state particles in their  $\gamma\gamma \rightarrow \omega J/\psi$  event sample [6]. The important angles for distinguishing  $J = 2^+$  from  $J = 0^\pm$  are  $\theta_n^*$ , the angle between  $\vec{n}$ , the normal to the  $\omega \rightarrow \pi^+\pi^-\pi^0$  decay plane, and the  $\gamma\gamma$  axis in the omega rest frame, and  $\theta_{in}$ , the angle between  $\vec{n}$  and the direction of the  $\ell^+$  from  $J/\psi \rightarrow \ell^+\ell^-$  decay (see Fig. 2a). Figure 2b shows the BaBar  $\cos\theta_n^*$  distribution together with the expectation for  $J = 0^\pm$  as a solid red line and  $J = 2^+$  as a dashed blue curve. There is a strong  $\chi^2$  penalty for the near-zero event likelihood near  $\cos\theta_n^* = \pm 1$  for the  $J = 2^+$  hypothesis to fluctuate *upward* to the observed levels of  $\sim 8$  and  $\sim 9$  events, and this is the main support BaBar's  $J = 0$  conclusion. The  $J = 2$  hypothesis seems to fit the BaBar  $\cos\theta_{in}$  distribution (see Fig. 2c) better than that for  $J = 0$ . But in this case, the likelihood of  $\sim 6$  expected events near  $\cos\theta_{in} = \pm 1$  to fluctuate *downward* to the observed  $\simeq 2$  events is not so improbable. With  $J = 0$  established, the  $0^+$  vs.  $0^-$  discrimination mostly relies on the angle  $\theta_n$ , which is the angle between the  $\omega$ 's flight path and  $\vec{n}$  in the  $\omega J/\psi$  restframe. The BaBar  $\cos\theta_n$  distribution shown in Fig. 2d favors  $0^+$  over  $0^-$ , mostly because of the  $\simeq 10$  events near  $\cos\theta_n = +1$ , where the  $0^-$  expectation is zero.

Babar's  $J^{PC} = 0^{++}$  assignment led them to suggest it as a suitable candidate for the  $2^3P_0$  charmonium state, commonly known as the  $\chi'_{c0}$ , and it was listed as such in the 2014 PDG tables [8]. However, this assignment had some problems and was challenged for a number of reasons [9]: the partial width for  $X(3915) \rightarrow \omega J/\psi$ , which would be an OZI-suppressed decay mode for a charmonium state, was too large; the lack of evidence for  $X(3915) \rightarrow D\bar{D}$ , which would be the dominant mode for the  $\chi'_{c0}$ ; and the small,  $\simeq 9$  MeV, mass splitting between the  $\chi'_{c2}$  and the  $X(3915)$ , which is an order-of-magnitude lower than the smallest theoretical estimates for  $M_{\chi'_{c2}} - M_{\chi'_{c0}}$  [10, 11]. This assignment was finally put to rest in 2017 by Belle, when they reported the observation of the  $X^*(3860)$ , a  $D\bar{D}$  resonance with mass  $3862^{+47}_{-35}$  MeV in  $e^+e^- \rightarrow J/\psi D\bar{D}$  annihilations with preferred spin-parity of  $J^{PC} = 0^{++}$  [12]. These properties, particularly the strong  $D\bar{D}$  decay mode, match well the expectations for the  $\chi'_{c0}$ , and the  $X^*(3862)$  is clearly a much stronger candidate for this state than the  $X(3915)$ .



**Figure 2.** a) Directions used in the BaBar study of  $\gamma\gamma \rightarrow \omega J/\psi$ , where  $\omega \rightarrow \pi^+\pi^-\pi^0$  and  $J/\psi \rightarrow \ell^+\ell^-$ . b) Comparison of the  $\cos\theta_n^*$  distribution with  $J^P = 0^\pm$  (solid red) and  $2^+$  (dashed blue) expectations. c) The corresponding plot for  $\cos\theta_{in}$ . d) The  $\cos\theta_n$  distribution with expectations for  $0^+$  in solid red and  $0^-$  in dashed blue. (From ref. [6].)


### 3 Is it the $\chi'_{c2}$ charmonium state?

The  $\chi'_{c2}$  was first spotted by Belle [13] and subsequently confirmed by BaBar [14] as a prominent  $M(D\bar{D})$  peak in the two-photon fusion process  $\gamma\gamma \rightarrow D\bar{D}$  that has a distinct  $\sin^4\theta^*$  production angle dependence that is characteristic of a  $J = 2$  state. The mass and width [7]:

$$M(\chi'_{c2}) = 3927.2 \pm 2.6 \text{ MeV} \quad \text{and} \quad \Gamma(\chi'_{c2}) = 24.0 \pm 6.0 \text{ MeV}, \quad (4)$$

are consistent with charmonium expectations for the  $\chi'_{c2}$  and there are no reasons to question this assignment. The Belle (BaBar)  $M(D\bar{D})$  and  $dN/d|\cos\theta^*|$  distributions are shown in Fig. 3a (b). Belle and BaBar measurements of its two-photon production rate are also in good agreement and are characterized by the product

$$\Gamma_{\gamma\gamma}(\chi'_{c2}) \times \mathcal{B}(\chi'_{c2} \rightarrow D\bar{D}) = 210 \pm 40 \text{ eV}. \quad (5)$$




**Figure 3.** a) *left:* The  $M(D\bar{D})$  distribution for  $\gamma\gamma \rightarrow D\bar{D}$ . The open histogram the  $D$  mass-sideband-determined background. The solid (dashed) curve shows results of a fit that includes (excludes) a  $\chi'_{c2}$  signal. *right:*  $dN/d|\cos\theta^*|$  for peak-region events with a solid (dashed) curve showing  $J = 2$  ( $J = 0$ ) expectations. The histogram is the non-resonant contribution. (From ref. [13].) b) Corresponding plots from BaBar [14].

BaBar's  $J^{PC} = 0^{++}$  assignment for the  $X(3915)$  was based on a comparison to a  $2^{++}$  scenario that only considered a helicity-2 component ( $h_2$ ) and ignored the possibility of any helicity-0 contribution. This assumption of “helicity-2 dominance” originate from a theoretical analysis that found that in two-photon production of tensor mesons, the helicity-0 component ( $h_0$ ) is zero in the non-relativistic limit [15]. The authors of ref. [16] point out that in the case of charmonium, the suppression of helicity-0 contributions only applies to mesons that are 100%  $c\bar{c}$ , which is generally considered to be unlikely for charmonium mesons with masses above the  $2m_D$  open-charm threshold (see, e.g., ref. [17]).

This is important because if the  $J^{PC}$  of the  $X(3915)$  is  $2^{++}$ , the mass peak identified with the  $X(3915)$  could be conceivably be due to an  $\omega J/\psi$  decay mode of the  $\chi_{c2}(2P)$  charmonium

state. The dashed lines in Fig. 4a show the ref. [16] comparison of the Belle  $M(D\bar{D})$  and  $|\cos\theta^*|$  with an  $h_0 \simeq 1.5h_2$  mixture to represent the  $X(3915)$ . Figure 4b) shows BaBar’s  $\cos\theta_n^*$  and  $\cos\theta_{in}$  distributions with expectations for  $0^{++}$ , and  $2^{++}$  with  $h = 0$  &  $h = 2$ . With the inclusion of some  $h = 0$  contribution, the  $\chi^2$  distinction between  $0^{++}$  and  $2^{++}$  angular distributions is diminished and the authors conclude that the  $X(3915)$  could be a  $\chi'_{c2}$  state that contains a sizable non- $c\bar{c}$  component.



**Figure 4.** a) Belle  $M(D\bar{D})$  (left) and  $|\cos\theta^*|$  (right) distributions for  $\gamma\gamma \rightarrow D\bar{D}$  production. The solid (dashed) curves show expectations for  $h_0 = 0$  ( $h_0 = 1.5h_2$ ). b) BaBar  $\cos\theta_n^*$  distribution (left) with a solid (dotted) curve showing expectations for  $2^{++}$  with  $h = 0$  ( $h = 2$ ); the dashed curve is for  $0^{++}$ . (right) The  $\cos\theta_{in}$  distribution with a solid curve for  $2^{++}$  with  $h = 0$  or 2, and a dashed curve for  $0^{++}$ . (From ref. [16].)

### 3.1 Other aspects of the $X(3915) = \chi'_{c2}$ assignment

In addition to violating helicity-2 dominance, which ref. [16] claims may not be a problem, there are other concerns with the  $X(3915) = \chi'_{c2}$  assignment. These are briefly discussed here.

#### 3.1.1 Mass and width differences

Belle and BaBar measurements of the  $\gamma\gamma \rightarrow \omega J/\psi$  mass peak,  $3915 \pm 4$  and  $3919 \pm 3$  MeV, respectively, are both lower, by  $\simeq 2\sigma$ , than their respective  $\chi'_{c2} \rightarrow D\bar{D}$  mass peak measurements,  $3929 \pm 5$  and  $3927 \pm 3$  MeV. Since the measurements reference well known masses –  $\omega$  and  $J/\psi$  for the  $X(3915)$  and  $D$ -meson for the  $\chi'_{c2}$  – systematic effects are small.

On the other hand, a recent LHCb report on the  $M(D\bar{D})$  distribution for inclusive  $D$ -meson pair production in high energy proton-proton collisions included observation of a distinct peak in the  $\chi'_{c2}$  mass region, shown in Fig. 5a, with mass  $M = 3921.9 \pm 0.6 \pm 0.2$  MeV,  $2\sigma$  below the  $\chi'_{c2}$  value listed in eqn. 4 [18]. The reported width,  $\Gamma = 36.6 \pm 1.9 \pm 0.9$  MeV, is  $2\sigma$  higher than the eqn. 4 value. The LHCb group attributes this peak to the  $\chi'_{c2}$ .


Figure 5b shows recent BESIII  $M(\omega J/\psi)$  results for  $e^+e^- \rightarrow Y(4220) \rightarrow \gamma\omega J/\psi$ , where there is a strong  $X(3872) \rightarrow \omega J/\psi$  signal and  $3\sigma$  “evidence” for two higher mass peaks [19]. The fitted mass of the middle peak is  $M = 3926.4 \pm 2.5$  MeV, near the Belle and BaBar results for  $\chi'_{c2} \rightarrow D\bar{D}$ . Thus, the current situation with mass measurements is inconclusive.

#### 3.1.2 A large OZI-violating $\omega J/\psi$ decay width for a $[c\bar{c}]$ meson

With the  $\Gamma_{\gamma\gamma} \times \mathcal{B}$  values listed in eqns. 3 and 5, the  $\chi'_{c2}$  assignment implies that

$$\frac{\mathcal{B}(\chi'_{c2} \rightarrow \omega J/\psi)}{\mathcal{B}(\chi'_{c2} \rightarrow D\bar{D})} = 0.05 \pm 0.02, \quad (6)$$

which is large for an OZI-rule-violating decay of an above-open-charm-threshold charmonium state, and more than an order-of-magnitude higher than the measured corresponding



**Figure 5.** a) The  $M(D^+D^-)$  distribution for inclusive  $D$ -meson pair production at the LHCb. The peak at 3842 MeV is the first observation of the  $\psi_3$ , the  $1^3D_3$  charmonium level. The broader peak near 3920 MeV is attributed by the LHCb group to the  $\chi'_{c2}$  [18]. b) The  $M(\omega J/\psi)$  distribution for  $e^+e^- \rightarrow Y(4220) \rightarrow \omega J/\psi$  events from BESIII. An  $X(3872) \rightarrow \omega J/\psi$  signal is evident. Additional peaks near 3925 MeV and 3960 MeV each have about  $3\sigma$  significance [19]. c)  $B^+ \rightarrow K^+\chi_{c1}$  and  $K^+\chi_{c2}$  signals from the full Belle data set [20].

ratio for  $\psi'' \rightarrow \pi^+\pi^- J/\psi$  and  $D\bar{D}$ . If  $\chi'_{c2} \rightarrow D\bar{D}$  and  $D\bar{D}^*$  are the dominant decay modes and  $\Gamma_{\chi'_{c2}}(D\bar{D}^*) \approx \Gamma_{\chi'_{c2}}(D\bar{D})$  (as predicted in ref. [21]), then  $\Gamma_{\chi'_{c2}}(\omega J/\psi) > 200$  keV (at the  $\sim 90\%$  CL), and much larger than any measured OZI-violating width for a charmonium state.

### 3.1.3 $\mathcal{B}(B \rightarrow K\chi'_{c2}) \gg \mathcal{B}(B \rightarrow K\chi_{c2})$ ?

In 2011, with their full event sample accumulated over ten years, Belle reported  $\sim 3\sigma$  evidence for  $B^+ \rightarrow K^+\chi_{c2}$  based on the  $33 \pm 11$  event signal shown in Fig. 5c [20]. The inferred branching fraction,  $\mathcal{B}(B^+ \rightarrow K^+\chi_{c2}) = 1.1 \pm 0.4 \times 10^{-5}$ , is smaller than the *product* branching fraction for  $X(3915) \rightarrow \omega J/\psi$  production in  $B^+$  meson decays (eqn. 2). Since  $\mathcal{B}(\chi'_{c2} \rightarrow D\bar{D})$  cannot exceed unity, eqn. 6 implies  $\mathcal{B}(\chi'_{c2} \rightarrow \omega J/\psi) < 0.08$  (90% CL). Thus, if the  $X(3915)$  produced in  $B \rightarrow K\omega J/\psi$  is the  $\chi'_{c2}$ , the  $B$ -meson decay width to  $K^+\chi'_{c2}$  would be more than an order of magnitude larger than that to  $K^+\chi_{c2}$ . This contradicts theoretical expectations that  $B \rightarrow K[c\bar{c}]$  decay widths decrease with increasing radial  $[c\bar{c}]$  quantum numbers [22].

Suppression of  $B \rightarrow K\chi_{c2}^{(\prime)}$  is not unexpected. The primary mechanism for  $B$ -meson ( $\bar{b}q$ ) decays to  $K[c\bar{c}]$  final states is  $\bar{b} \rightarrow \bar{c}$  plus a virtual  $W^+$  that, in turn, materializes as  $c\bar{s}$ . The final-state  $c$ - and  $\bar{c}$ -quark form the  $[c\bar{c}]$  state and the  $\bar{s}$ - and “spectator”  $q$ -quark form the  $K$ . This process is only allowed for  $J^{PC} = 0^+, 1^{--}$  and  $1^{++}$   $[c\bar{c}]$  states, decays to  $[c\bar{c}]$  states with other  $J^{PC}$  values are higher-order and expected to be “factorization suppressed” [23]. The Belle results on  $B \rightarrow K\chi_{c2}$  shown in Fig. 5c demonstrate that for  $J^{PC} = 2^{++}$   $[c\bar{c}]$  states, factorization suppression is very effective:  $\mathcal{B}(B \rightarrow K\chi_{c2}) < 0.04 \times \mathcal{B}(B \rightarrow K\chi_{c1})$  (90% CL).

## 4 Summary and conclusions

Despite its observation by different experiments in a variety of production channels, the nature of the  $X(3915)$  remains a mystery. If it is a nonstandard  $XYZ$  meson, it cannot be easily interpreted by any of the proposed models for these states. For example: its mass is too low for a QCD-hybrid [24], and not near an appropriate threshold for a molecular state or a cusp effect [25]; the lack of evidence for a  $\eta\eta_c$  decay mode [26] is problematic for a diquark-diantiquark assignment [27]. Thus, if it is an  $XYZ$  meson, it is a very interesting one.

The sum total of existing data on  $\omega J/\psi$  and  $D\bar{D}$  production in the  $\sim 3925$  MeV mass region *cannot* be explained as being simply due to the  $\chi'_{c2}$  charmonium state. While a (tenuous) case could be made that the near-3925 MeV mass peaks seen by the LHCb in  $pp \rightarrow D\bar{D}X$ , Belle and BaBar in  $\gamma\gamma \rightarrow \omega J/\psi$  &  $D\bar{D}$  and BESIII in  $Y(4220) \rightarrow \gamma\omega J/\psi$  are all due to decays of the  $\chi'_{c2}$ , the existing evidence is not conclusive. Moreover, a very strong case can be made *against* a  $\chi'_{c2}$  interpretation of the  $\omega J/\psi$  peak seen in  $B \rightarrow K\omega J/\psi$  decays.

More refined mass and width measurements are needed, and reliable, separate  $J^{PC}$  determinations for the  $\omega J/\psi$  peaks produced via  $\gamma\gamma$  fusion, radiative  $Y(4220)$  transitions, and  $B$ -meson decays that eschew the helicity-2 dominance constraint are essential. The LHCb group has demonstrated that they can isolate clean  $B^+ \rightarrow K^+ \omega J/\psi$  signals with good efficiency [28] and I look forward to high-statistics results from them in the near future.

## 5 Acknowledgements

I congratulate Phi-to-Psi-2018 organizers for an interesting and provocative meeting. This work is supported by the CAS President's International Fellowship Initiative.

## References

- [1] S.L. Olsen, T. Skwarnicki, D. Zieminska, Rev. Mod. Phys. **90**, 015003 (2018), 1708.04012
- [2] K. Abe et al. (Belle), Phys. Rev. Lett. **94**, 182002 (2005), hep-ex/0408126
- [3] B. Aubert et al. (BaBar), Phys. Rev. Lett. **101**, 082001 (2008), 0711.2047
- [4] P. del Amo Sanchez et al. (BaBar), Phys. Rev. **D82**, 011101 (2010), 1005.5190
- [5] S. Uehara et al. (Belle), Phys. Rev. Lett. **104**, 092001 (2010), 0912.4451
- [6] J.P. Lees et al. (BaBar), Phys. Rev. **D86**, 072002 (2012), 1207.2651
- [7] M. Tanabashi et al. (Particle Data Group), Phys. Rev. **D98**, 030001 (2018)
- [8] K.A. Olive et al. (Particle Data Group), Chin. Phys. **C38**, 090001 (2014)
- [9] F.K. Guo, U.G. Meissner, Phys. Rev. **D86**, 091501 (2012), 1208.1134
- [10] H. Wang, Y. Yang, J. Ping, Eur. Phys. J. **A50**, 76 (2014)
- [11] S.L. Olsen, Phys. Rev. **D91**, 057501 (2015), 1410.6534
- [12] K. Chilikin et al. (Belle), Phys. Rev. **D95**, 112003 (2017), 1704.01872
- [13] S. Uehara et al. (Belle), Phys. Rev. Lett. **96**, 082003 (2006), hep-ex/0512035
- [14] B. Aubert et al. (BaBar), Phys. Rev. **D81**, 092003 (2010), 1002.0281
- [15] M. Krammer, H. Krasemann, Phys. Lett. **73B**, 58 (1978)
- [16] Z.Y. Zhou, Z. Xiao, H.Q. Zhou, Phys. Rev. Lett. **115**, 022001 (2015), 1501.00879
- [17] M.R. Pennington, D.J. Wilson, Phys. Rev. **D76**, 077502 (2007), 0704.3384
- [18] R. Aaij et al. (LHCb) (2019), 1903.12240
- [19] M. Ablikim et al. (BESIII) (2019), 1903.04695
- [20] V. Bhardwaj et al. (Belle), Phys. Rev. Lett. **107**, 091803 (2011), 1105.0177
- [21] T. Barnes, S. Godfrey, E.S. Swanson, Phys. Rev. **D72**, 054026 (2005), hep-ph/0505002
- [22] G.T. Bodwin et al., Phys. Rev. **D46**, R3703 (1992), hep-ph/9208254
- [23] M. Beneke et al., Phys. Rev. Lett. **83**, 1914 (1999), hep-ph/9905312
- [24] L. Liu et al. (Hadron Spectrum), JHEP **07**, 126 (2012), 1204.5425
- [25] S.L. Olsen, EPJ Web Conf. **202**, 01003 (2019), 1812.10947
- [26] A. Vinokurova et al. (Belle), JHEP **06**, 132 (2015), [Erratum: JHEP02,088(2017)], 1501.06351
- [27] R.F. Lebed, A.D. Polosa, Phys. Rev. **D93**, 094024 (2016), 1602.08421
- [28] G. Andreassi, Ph.D. thesis, University of Rome, Sapienza (2014)